

thm_2EConseqConv_2EAND__CLAUSES__FX (TMR1AxUoopT3dZicb8xsFd1t8xtsqvLyvYY)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2))) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x)$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A_27a}))))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((\text{True} \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \wedge \text{True}) \Leftrightarrow \\ & (p \ V0t)) \wedge (((\text{False} \wedge (p \ V0t)) \Leftrightarrow \text{False}) \wedge (((p \ V0t) \wedge \text{False}) \Leftrightarrow \text{False}) \wedge \\ & (((p \ V0t) \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall A_27a. \text{nonempty } A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1Q \in \\ & (2^{A_27a}). ((\forall V2x \in A_27a. ((p \ (\text{ap } V0P \ V2x)) \wedge (p \ (\text{ap } V1Q \ V2x)))) \Leftrightarrow \\ & ((\forall V3x \in A_27a. (p \ (\text{ap } V0P \ V3x))) \wedge (\forall V4x \in A_27a. (p \ (\text{ap } V1Q \ V4x))))))))) \end{aligned} \quad (2)$$

Theorem 1 $(\forall V0t \in 2. ((\text{False} \wedge (p \ V0t)) \Leftrightarrow \text{False}))$.