

thm_2EConseqConv_2EAND__CLAUSES__XX
(TMYQS7idcPJMyd7wPFTSBCwmqkKKFTqAmvZ)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V1Q \in (2^{A-27a}).(\lambda V2x \in A_27a.((p (ap V0P V2x)) \wedge (p (ap V1Q V2x)))) \Leftrightarrow ((\lambda V3x \in A_27a.(p (ap V0P V3x))) \wedge (\lambda V4x \in A_27a.(p (ap V1Q V4x))))))))))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0P \in (2^{A-27a}).(\forall V1Q \in \\ & (2^{A-27a}).((\forall V2x \in A_27a.((p (ap V0P V2x)) \wedge (p (ap V1Q V2x)))) \Leftrightarrow \\ & ((\forall V3x \in A_27a.(p (ap V0P V3x))) \wedge (\forall V4x \in A_27a.(p (ap V1Q V4x))))))))) \end{aligned} \quad (2)$$

Theorem 1 $(\forall V0t \in 2.(((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))$.