

thm_2EConseqConv_2Eforall_eq_thm (TMKNG-
BKitqFtjcWfT4FAMJyw2A6D7UiuWPf)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{2}$$

Theorem 1

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0P \in (2^{A_27a}).(\forall V1Q \in (2^{A_27a}).((\forall V2s \in A_27a.((p (ap V0P V2s)) \Leftrightarrow (p (ap V1Q V2s)))) \Rightarrow ((\forall V3s \in A_27a.(p (ap V0P V3s))) \Leftrightarrow (\forall V4s \in A_27a.(p (ap V1Q V4s))))))))))$$