

thm_2EEncodeVar_2Eof_length_exists_zero (TMZ88pN8aJPjbouvktmQsytzXS7k2AA9Beo)

October 26, 2020

Let $ty_2Elist_2Elist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Elist_2Elist\ A0) \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (2)$$

Let $c_2EEncodeVar_2Eof_length : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2EEncodeVar_2Eof_length\ A_27a \in ((2^{(ty_2Elist_2Elist\ A_27a)})_{ty_2Enum_2Enum}) \quad (3)$$

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_2IN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap\ V1f\ V0x)))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A_27a})))$

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t)))$

Definition 7 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if\ (\exists x \in A.p\ (ap\ P\ x))\ \mathbf{then}\ (the\ (\lambda x.x \in A \wedge p\ x))$ of type $\iota \Rightarrow \iota$.

Definition 8 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ V0P\ (ap\ (c_2Emin_2E_40\ A_27a)\ P)))$

Definition 9 We define $c_2Ebool_2ERES_EXISTS$ to be $\lambda A_27a : \iota.(\lambda V0p \in (2^{A_27a}).(\lambda V1m \in (2^{A_27a}).(ap\ V1m\ p)))$

Let $c_2Elist_2ENIL : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Elist_2ENIL\ A_27a \in (ty_2Elist_2Elist\ A_27a) \quad (4)$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (5)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (6)$$

Definition 10 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Elist_2ELENGTH : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Elist_2ELENGTH\ A_27a \in (ty_2Enum_2Enum^{(ty_2Elist_2Elist\ A_27a)}) \quad (7)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0l \in (ty_2Elist_2Elist\ A_27a). (\forall V1n \in ty_2Enum_2Enum. ((p\ (ap\ (ap\ (c_2Ebool_2EIN \\ (ty_2Elist_2Elist\ A_27a))\ V0l)\ (ap\ (c_2EEncodeVar_2Eof_length\ A_27a)\ V1n))) \Leftrightarrow ((ap\ (c_2Elist_2ELENGTH\ A_27a)\ V0l) = V1n)))) \end{aligned} \quad (8)$$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (10)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a. (p\ V0t) \Leftrightarrow (p\ V0t))) \quad (11)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1a \in A_27a. ((\exists V2x \in A_27a. ((V2x = V1a) \wedge (p\ (ap\ V0P\ V2x)))) \Leftrightarrow (p\ (ap\ V0P\ V1a)))))) \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0P \in (2^{A.27a}). (\forall V1f \in \\ & (2^{A.27a}). ((p\ (ap\ (ap\ (c.2Ebool.2ERES_EXISTS\ A.27a)\ V0P)\ V1f))) \Leftrightarrow \\ & (\exists V2x \in A.27a. ((p\ (ap\ (ap\ (c.2Ebool.2EIN\ A.27a)\ V2x)\ V0P)) \wedge \\ & (p\ (ap\ V1f\ V2x)))))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0l \in (ty.2Elist.2Elist \\ & A.27a). ((ap\ (c.2Elist.2ELENGTH\ A.27a)\ V0l) = c.2Enum.2E0) \Leftrightarrow (\\ & V0l = (c.2Elist.2ENIL\ A.27a))) \end{aligned} \quad (15)$$

Theorem 1

$$\begin{aligned} & \forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0phi \in (2^{(ty.2Elist.2Elist\ A.27a)}). \\ & ((p\ (ap\ (ap\ (c.2Ebool.2ERES_EXISTS\ (ty.2Elist.2Elist\ A.27a)) \\ & (ap\ (c.2EEncodeVar.2Eof_length\ A.27a)\ c.2Enum.2E0))\ (\lambda V1w \in \\ & (ty.2Elist.2Elist\ A.27a). (ap\ V0phi\ V1w)))) \Leftrightarrow (p\ (ap\ V0phi\ (c.2Elist.2ENIL \\ & A.27a)))) \end{aligned}$$