

thm\_2EEncode\_2Ewf\_encode\_bnum  
 (TMXyjqvnMzBX-  
 hZFQwqkJF8NAGMRHtyEHcwn)

October 26, 2020

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (1)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$\text{nonempty } ty\_2Enum\_2Enum \quad (2)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (3)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 3** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (4)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (5)$$

**Definition 4** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 5** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A\_27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x)))$

**Definition 6** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. (ap\ c\_2Enum\_2EABS\_num\ m)$

Let  $c_2$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (6)$$

**Definition 7** We define  $c\_2Earithmetic\_2EBIT2$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmetic\_2EBIT2$

**Definition 8** We define  $c_2\text{Earthmetic\_ENUMERAL}$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Let  $c_2 \in \text{arithmic\_EXP} : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2EEEXP \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \\ (7)$$

**Definition 9** We define  $c\_2Ebool\_2EF$  to be  $(ap\ (c\_2Ebool\_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 10** We define  $c_2 \in \text{min}_2 \cup \text{max}_2$  to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o} (p \rightarrow_p Q)$  of type  $\iota$ .

**Definition 11** We define  $c_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E))$

**Definition 12** We define  $c_{\text{Ebool}} : \mathbb{E} \rightarrow \mathbb{F}_{\text{5C}}$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap(c_{\text{Ebool}} \_2E\_21) 2)) (\lambda V2t \in$

**Definition 13** We define  $c_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p \text{ (ap } P \text{ } x)) \text{ then } (\lambda x.x \in A \wedge$  of type  $\iota \Rightarrow \iota$ .

**Definition 14** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.\_27a : \iota.(\lambda V0P \in (2^A\_{-27}a)).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 15** We define  $c_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 16** We define  $c\_2EEncode\_2Ewf\_pred$  to be  $\lambda A\_27a : \iota. \lambda V0p \in (2^A\_{27a}).(ap (c\_2Ebool\_2E\_3F A))$

**Definition 17** We define  $c \in \text{Encode}_{\text{Ewf\_pred\_bnum}}$  to be  $\lambda V0m \in \text{ty\_Enum}.\lambda V1p \in (2^{\text{ty\_Enum}})$

Let  $c_2$  be given. Assume the following.

$c\_2Earithmetic\_2EMOD \in ((ty\_2Enum\_2Enum^{ty\_2Enum^{ty\_2Enum\_2Enum}})^*$

(8)

Let  $t : \text{list } \text{list} \rightarrow \text{list}$  be given. Assume the following.

$\forall A0 \rightarrow nonempty, A0 \Rightarrow nonempty, (tu : 2El) \in 2El$

needs  $2E$  needs,  $b_{\text{num}}$  to be given. Assume the following:

$$2E_E - 1 - 2E_U = 1 - 1 \quad \in ((1 - 2E_U) + 2E_U) + 2tu/2E_{num}/2E_{nu}$$

(10)

Let  $c\_2Elist\_2EisPREFIX : \iota \rightarrow \iota$  be given. Assume the following.

$$V_{\text{TE}}(t) \approx \frac{1}{2} \left( \alpha_1 + \alpha_2 \right) t^2 + \frac{1}{2} \left( \alpha_1 - \alpha_2 \right) \cos(\omega_0 t) \quad (11)$$

**Definition 19** We define  $c\_2EEncode\_2Ewf\_encoder$  to be  $\lambda A\_27a : \iota. \lambda V0p \in (2^{A\_27a}). \lambda V1e \in ((ty\_2Elist \_2Ewf \_2Ewf) \_2Ewf \_2Ewf \_2Ewf) . V1e$

**Definition 20** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1p \in (2^{ty\_2Enum\_2Enum})). \\ & ((p (ap (ap c\_2EEncode\_2Ewf\_pred\_bnum V0m) V1p)) \Rightarrow (p (ap (ap c\_2EEncode\_2Ecollision\_free \\ & V0m) V1p)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1p \in (2^{ty\_2Enum\_2Enum})). \\ & ((p (ap (ap (c\_2EEncode\_2Ewf\_encoder ty\_2Enum\_2Enum) V1p) (ap \\ & c\_2EEncode\_2Eencode\_bnum V0m))) \Leftrightarrow (p (ap (ap c\_2EEncode\_2Ecollision\_free \\ & V0m) V1p)))))) \end{aligned} \quad (13)$$

Assume the following.

$$True \quad (14)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p \\ & V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (15)$$

Assume the following.

$$(\forall V0t \in 2. (((p V0t) \Rightarrow False) \Rightarrow (\neg(p V0t)))) \quad (16)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(p V0t)) \Rightarrow ((p V0t) \Rightarrow False))) \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (( \\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t))))))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & ((\forall V0t \in 2. ((\neg(\neg(p V0t)) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ & ((\neg False) \Leftrightarrow True))) \end{aligned} \quad (19)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg \\ & (p V0t))))))) \end{aligned} \quad (20)$$

Assume the following.

$$\forall A_27a. \text{nonempty } A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). ((\neg(\forall V1x \in A_27a. (p (ap V0P V1x)))) \Leftrightarrow (\exists V2x \in A_27a. (\neg(p (ap V0P V2x))))) \quad (21)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((p V0A) \vee (p V1B)) \Leftrightarrow ((p V1B) \vee (p V0A)))) \quad (22)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (23)$$

Assume the following.

$$(\forall V0A \in 2. ((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow \text{False}))) \quad (24)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((p V0A) \vee (p V1B))) \Rightarrow \text{False}) \Leftrightarrow ((p V0A) \Rightarrow \text{False}) \Rightarrow ((\neg(p V1B)) \Rightarrow \text{False}))) \quad (25)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow \text{False}) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow \text{False}))) \quad (26)$$

Assume the following.

$$(\forall V0A \in 2. (((\neg(p V0A)) \Rightarrow \text{False}) \Rightarrow (((p V0A) \Rightarrow \text{False}) \Rightarrow \text{False}))) \quad (27)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow (p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee ((\neg(p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p))))))))))) \quad (28)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow (p V1q) \wedge (p V2r))) \Leftrightarrow (((p V0p) \vee ((\neg(p V1q)) \vee (\neg(p V2r)))) \wedge (((p V1q) \vee (\neg(p V0p))) \wedge ((p V2r) \vee (\neg(p V0p)))))))) \quad (29)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow (p V1q) \vee (p V2r))) \Leftrightarrow (((p V0p) \vee ((\neg(p V1q)) \wedge (\neg(p V2r)))) \wedge (((p V0p) \vee ((\neg(p V2r)) \wedge (\neg(p V1q)))) \wedge (((p V1q) \vee ((p V2r) \vee (\neg(p V0p))))))))))) \quad (30)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow ((p V1q) \Rightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((\neg(p V1q)) \vee ((p V2r) \vee (\neg(p V0p)))))))))) \quad (31)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p))))))) \quad (32)$$

### Theorem 1

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1p \in (2^{ty\_2Enum\_2Enum}). \\ & ((p (ap (ap c\_2EEncode\_2Ewf\_pred\_bnum V0m) V1p)) \Rightarrow (p (ap (ap ( \\ & c\_2EEncode\_2Ewf\_encoder ty\_2Enum\_2Enum) V1p) (ap c\_2EEncode\_2Eencode\_bnum \\ & V0m))))))) \end{aligned}$$