

thm_2EEncode_2Ewf__encode__option (TMQs8gtg1xUhWVPQk4WHsP6MnkpprGMoeSD)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Let $ty_2Elist_2Elist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Elist_2Elist\ A0) \quad (1)$$

Let $c_2Elist_2EisPREFIX : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Elist_2EisPREFIX\ A_27a \in ((2^{(ty_2Elist_2Elist\ A_27a)})^{(ty_2Elist_2Elist\ A_27a)}) \quad (2)$$

Definition 2 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 3 We define $c_2Ebool_2E_2$ to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A_27a}))\ (\lambda V1Q \in 2.V1Q))\ (\lambda V1R \in 2.V1R)))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))))$

Definition 6 We define $c_2EEncode_2Ewf_encoder$ to be $\lambda A_27a : \iota.\lambda V0p \in (2^{A_27a}).\lambda V1e \in ((ty_2Elist_2Elist\ 2)^{A_27a})$

Let $ty_2Eoption_2Eoption : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Eoption_2Eoption\ A0) \quad (3)$$

Let $c_2EEncode_2Eencode_option : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2EEncode_2Eencode_option\ A_27a \in (((ty_2Elist_2Elist\ 2)^{(ty_2Eoption_2Eoption\ A_27a)})^{((ty_2Elist_2Elist\ 2)^{A_27a})}) \quad (4)$$

Let $c_2Eoption_2Eoption_CASE : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Eoption_2Eoption_CASE\ A_27a\ A_27b \in (((A_27b^{(A_27b^{A_27a})})^{A_27b})^{(ty_2Eoption_2Eoption\ A_27a)}) \quad (5)$$

Definition 7 We define $c_2EEncode_2Elift_option$ to be $\lambda A_27a : \iota. \lambda V0p \in (2^{A-27a}). \lambda V1x \in (ty_2Eoption_...$

Definition 8 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21\ 2) (\lambda V0t \in 2.V0t))$.

Definition 9 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E\ V0t) c_2Ebool_2EF$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A. \lambda P \in 2^A. \mathbf{if} (\exists x \in A. p (ap\ P\ x)) \mathbf{then}$ (the $(\lambda x. x \in A \wedge$
of type $\iota \Rightarrow \iota$).

Definition 11 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap\ V0P\ (ap\ (c_2Emin_2E_40$

Definition 12 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21\ 2) (\lambda V2t \in$

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \tag{6}$$

Definition 13 We define c_2Eone_2Eone to be $(ap (c_2Emin_2E_40\ ty_2Eone_2Eone) (\lambda V0x \in ty_2Eone_2Eone$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. nonempty\ A0 \Rightarrow \forall A1. nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \tag{7}$$

Let $c_2Esum_2EABS_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow \forall A_27b. nonempty\ A_27b \Rightarrow c_2Esum_2EABS_sum\ A_27a\ A_27b \in ((ty_2Esum_2Esum\ A_27a\ A_27b)^{((2^{A-27b})^{A-27a})^2}) \tag{8}$$

Definition 14 We define c_2Esum_2EINR to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27b. (ap (c_2Esum_2EABS$

Let $c_2Eoption_2Eoption_ABS : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow c_2Eoption_2Eoption_ABS\ A_27a \in ((ty_2Eoption_2Eoption\ A_27a)^{(ty_2Esum_2Esum\ A_27a\ ty_2Eone_2Eone)}) \tag{9}$$

Definition 15 We define $c_2Eoption_2ENONE$ to be $\lambda A_27a : \iota. (ap (c_2Eoption_2Eoption_ABS\ A_27a) ($

Definition 16 We define c_2Esum_2EINL to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27a. (ap (c_2Esum_2EABS$

Definition 17 We define $c_2Eoption_2ESOME$ to be $\lambda A_27a : \iota. \lambda V0x \in A_27a. (ap (c_2Eoption_2Eoption$

Let $c_2Elist_2ECONS : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow c_2Elist_2ECONS\ A_27a \in (((ty_2Elist_2Elist\ A_27a)^{(ty_2Elist_2Elist\ A_27a)})^{A_27a}) \tag{10}$$

Let $c_2Elist_2ENIL : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Elist_2ENIL\ A_27a \in (ty_2Elist_2Elist\ A_27a) \quad (11)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow & ((\forall V0xb \in ((ty_2Elist_2Elist\ 2)^{A_27a}).((ap\ (ap\ (c_2EEncode_2Eencode_option\ A_27a)\ V0xb) \\ & (c_2Eoption_2ENONE\ A_27a)) = (ap\ (ap\ (c_2Elist_2ECONS\ 2)\ c_2Ebool_2EF \\ & (c_2Elist_2ENIL\ 2)))))) \wedge (\forall V1xb \in ((ty_2Elist_2Elist\ 2)^{A_27a}). \\ & (\forall V2x \in A_27a.((ap\ (ap\ (c_2EEncode_2Eencode_option\ A_27a) \\ & V1xb)\ (ap\ (c_2Eoption_2ESOME\ A_27a)\ V2x)) = (ap\ (ap\ (c_2Elist_2ECONS\ 2)\ c_2Ebool_2ET)\ (ap\ V1xb\ V2x)))))) \end{aligned} \quad (12)$$

Assume the following.

$$True \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (14)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} ((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ ((\neg False) \Leftrightarrow True))) \end{aligned} \quad (16)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (17)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (18)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg \\ p\ V0t)))))) \end{aligned} \quad (19)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow (p V1t2) \Rightarrow (p V2t3)) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (20)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1x_{.27} \in 2. (\forall V2y \in 2. (\forall V3y_{.27} \in 2. (((p V0x) \Leftrightarrow (p V1x_{.27})) \wedge ((p V1x_{.27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{.27})))))) \Rightarrow ((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{.27}) \Rightarrow (p V3y_{.27})))))) \quad (21)$$

Assume the following.

$$\forall A_{.27a}. \text{nonempty } A_{.27a} \Rightarrow (\forall V0opt \in (ty_2Eoption_2Eoption A_{.27a}). ((V0opt = (c_2Eoption_2ENONE A_{.27a})) \vee (\exists V1x \in A_{.27a}. (V0opt = (ap (c_2Eoption_2ESOME A_{.27a}) V1x)))))) \quad (22)$$

Assume the following.

$$\forall A_{.27a}. \text{nonempty } A_{.27a} \Rightarrow \forall A_{.27b}. \text{nonempty } A_{.27b} \Rightarrow ((\forall V0v \in A_{.27b}. (\forall V1f \in (A_{.27b}^{A_{.27a}}). ((ap (ap (ap (c_2Eoption_2Eoption_CASE A_{.27a} A_{.27b}) (c_2Eoption_2ENONE A_{.27a})) V0v) V1f) = V0v))) \wedge (\forall V2x \in A_{.27a}. (\forall V3v \in A_{.27b}. (\forall V4f \in (A_{.27b}^{A_{.27a}}). ((ap (ap (ap (c_2Eoption_2Eoption_CASE A_{.27a} A_{.27b}) (ap (c_2Eoption_2ESOME A_{.27a}) V2x)) V3v) V4f) = (ap V4f V2x))))))) \quad (23)$$

Assume the following.

$$\forall A_{.27a}. \text{nonempty } A_{.27a} \Rightarrow (\forall V0x \in A_{.27a}. (\forall V1y \in A_{.27a}. (((ap (c_2Eoption_2ESOME A_{.27a}) V0x) = (ap (c_2Eoption_2ESOME A_{.27a}) V1y)) \Leftrightarrow (V0x = V1y)))) \quad (24)$$

Assume the following.

$$\forall A_{.27a}. \text{nonempty } A_{.27a} \Rightarrow ((\forall V0l \in (ty_2Elist_2Elist A_{.27a}). ((p (ap (ap (c_2Elist_2EisPREFIX A_{.27a}) (c_2Elist_2ENIL A_{.27a})) V0l)) \Leftrightarrow True)) \wedge ((\forall V1x \in A_{.27a}. (\forall V2l \in (ty_2Elist_2Elist A_{.27a}). ((p (ap (ap (c_2Elist_2EisPREFIX A_{.27a}) (ap (ap (c_2Elist_2ECONS A_{.27a}) V1x) V2l)) (c_2Elist_2ENIL A_{.27a})) \Leftrightarrow False)))) \wedge (\forall V3x1 \in A_{.27a}. (\forall V4l1 \in (ty_2Elist_2Elist A_{.27a}). (\forall V5x2 \in A_{.27a}. (\forall V6l2 \in (ty_2Elist_2Elist A_{.27a}). ((p (ap (ap (c_2Elist_2EisPREFIX A_{.27a}) (ap (ap (c_2Elist_2ECONS A_{.27a}) V5x2) V6l2)) (ap (ap (c_2Elist_2ECONS A_{.27a}) V3x1) V4l1))) \Leftrightarrow ((V3x1 = V5x2) \wedge (p (ap (ap (c_2Elist_2EisPREFIX A_{.27a}) V6l2) V4l1)))))))))) \quad (25)$$

Theorem 1

$$\forall A_{.27a}. \text{nonempty } A_{.27a} \Rightarrow (\forall V0p \in (2^{A_{.27a}}). (\forall V1e \in ((ty_2Elist_2Elist 2)^{A_{.27a}}). ((p (ap (ap (c_2EEncode_2Ewf_encoder A_{.27a}) V0p) V1e)) \Rightarrow (p (ap (ap (c_2EEncode_2Ewf_encoder (ty_2Eoption_2Eoption A_{.27a}) (ap (c_2EEncode_2Elift_option A_{.27a}) V0p)) (ap (c_2EEncode_2Eencode_option A_{.27a}) V1e)))))) \quad (26)$$