

# thm\_2EEncode\_2Ewf\_\_encoder\_\_total (TMLR869qpt1U2e1A5ubopZpytaEd2uBPrPG)

October 26, 2020

Let  $ty\_2Elist\_2Elist : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Elist\_2Elist\ A0) \quad (1)$$

Let  $c\_2Elist\_2EisPREFIX : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Elist\_2EisPREFIX\ A\_27a \in ((2^{(ty\_2Elist\_2Elist\ A\_27a)})^{(ty\_2Elist\_2Elist\ A\_27a)}) \quad (2)$$

**Definition 1** We define  $c\_2Emin\_2E3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o\ (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Emin\_2E3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o\ (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .

**Definition 3** We define  $c\_2Ebool\_2E2T$  to be  $(ap\ (ap\ (c\_2Emin\_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ (ap\ (c\_2Emin\_2E3D\ (2^{A\_27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x)))$

**Definition 5** We define  $c\_2Ebool\_2E2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E21\ 2)\ (\lambda V2t \in 2.V2t)))$

**Definition 6** We define  $c\_2EEncode\_2Ewf\_encoder$  to be  $\lambda A\_27a : \iota.\lambda V0p \in (2^{A\_27a}).\lambda V1e \in ((ty\_2Elist\_2Elist\ A\_27a)^{A\_27a})$

**Definition 7** We define  $c\_2Ebool\_2E2F$  to be  $(ap\ (c\_2Ebool\_2E21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 8** We define  $c\_2Ebool\_2E2E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E2F))$

**Definition 9** We define  $c\_2Ecombin\_2EK$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.(\lambda V0x \in A\_27a.(\lambda V1y \in A\_27b.V0x))$

Assume the following.

$$True \quad (3)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p\ V0t) \Leftrightarrow (p\ V0t)))) \quad (4)$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \wedge True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \wedge (p \ V0t)) \Leftrightarrow False) \wedge (((p \ V0t) \wedge False) \Leftrightarrow False) \wedge \\
& (((p \ V0t) \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)))))) \quad (5)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Rightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Rightarrow True) \Leftrightarrow \\
& True) \wedge (((False \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (((p \ V0t) \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (( \\
& (p \ V0t) \Rightarrow False) \Leftrightarrow (\neg(p \ V0t)))))) \quad (6)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in \\
& A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x))) \quad (7)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow (\neg(p \ V0t))) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow (\neg( \\
& p \ V0t)))))) \quad (8)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p \ V0t1) \Rightarrow \\
& ((p \ V1t2) \Rightarrow (p \ V2t3))) \Leftrightarrow (((p \ V0t1) \wedge (p \ V1t2)) \Rightarrow (p \ V2t3)))))) \quad (9)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in 2. (\forall V1x\_27 \in 2. (\forall V2y \in 2. (\forall V3y\_27 \in \\
& 2. (((p \ V0x) \Leftrightarrow (p \ V1x\_27)) \wedge ((p \ V1x\_27) \Rightarrow ((p \ V2y) \Leftrightarrow (p \ V3y\_27)))) \Rightarrow \\
& (((p \ V0x) \Rightarrow (p \ V2y)) \Leftrightarrow ((p \ V1x\_27) \Rightarrow (p \ V3y\_27)))))) \quad (10)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall A\_27a.nonempty \ A\_27a \Rightarrow \forall A\_27b.nonempty \ A\_27b \Rightarrow ( \\
& \forall V0x \in A\_27a. (\forall V1y \in A\_27b. ((ap \ (ap \ (c\_2Ecombin\_2EK \\
& A\_27a \ A\_27b) \ V0x) \ V1y) = V0x)) \quad (11)
\end{aligned}$$

### Theorem 1

$$\begin{aligned}
& \forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0p \in (2^{A\_27a}). (\forall V1e \in \\
& ((ty\_2Elist\_2Elist \ 2)^{A\_27a}). ((p \ (ap \ (ap \ (c\_2EEncode\_2Ewf\_encoder \\
& A\_27a) \ (ap \ (c\_2Ecombin\_2EK \ 2 \ A\_27a) \ c\_2Ebool\_2ET)) \ V1e)) \Rightarrow (p \ ( \\
& ap \ (ap \ (c\_2EEncode\_2Ewf\_encoder \ A\_27a) \ V0p) \ V1e))))
\end{aligned}$$