

thm\_2EHolSmt\_2EF\_\_OR  
(TMdeEbeBFY88n86tMvA.JxWDeo7j4dU1kzG7)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Ebool_2ET` to be  $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define `c_2Ebool_2E_21` to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (P) (Q))$

**Definition 4** We define `c_2Ebool_2EF` to be  $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

**Definition 7** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

**Definition 8** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee \\ & (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p V0t)))))) \end{aligned} \tag{3}$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \tag{4}$$

Assume the following.

$$(\forall V0A \in 2.((p \ V0A) \Rightarrow ((\neg(p \ V0A)) \Rightarrow False))) \quad (5)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg(\neg(p \ V0A)) \vee (p \ V1B))) \Rightarrow False) \Leftrightarrow ((p \ V0A) \Rightarrow ((\neg(p \ V1B)) \Rightarrow False)))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p \ V0A)) \Rightarrow False) \Rightarrow ((p \ V0A) \Rightarrow False) \Rightarrow False)) \quad (7)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((p \ V0p) \Leftrightarrow (\neg(p \ V1q))) \Leftrightarrow (((p \ V0p) \vee (p \ V1q)) \wedge ((\neg(p \ V1q)) \vee (\neg(p \ V0p)))))) \quad (8)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p \ V0p) \Rightarrow (p \ V1q))) \Rightarrow (p \ V0p))) \quad (9)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p \ V0p) \Rightarrow (p \ V1q))) \Rightarrow (\neg(p \ V1q)))) \quad (10)$$

**Theorem 1**

$$(\forall V0p \in 2.(\forall V1q \in 2.(((False \vee (p \ V0p)) \Leftrightarrow (False \vee (p \ V1q))) \Rightarrow ((p \ V0p) \Leftrightarrow (p \ V1q))))$$