

thm_2EHolSmt_2EIMP__DISJ__2
(TMHZWmGJ6TMr4aTngSybWzYY3KzyMCWxErJ)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2))) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x)$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 7 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 8 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t)) (\text{c_2Ebool_2E_2F } 2)))$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((\text{True} \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow \text{True}) \Leftrightarrow \\ & (p \ V0t)) \wedge (((\text{False} \Leftrightarrow (p \ V0t)) \Leftrightarrow \neg(p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow \text{False}) \Leftrightarrow \neg(p \ V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p \ V0t))) \Leftrightarrow (p \ V0t))) \tag{3}$$

Assume the following.

$$(\forall V0A \in 2. ((p \ V0A) \Rightarrow ((\neg(p \ V0A)) \Rightarrow \text{False}))) \tag{4}$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p \ V0A) \vee (p \ V1B))) \Rightarrow False) \Leftrightarrow ((p \ V0A) \Rightarrow False) \Rightarrow ((\neg(p \ V1B)) \Rightarrow False)))) \quad (5)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p \ V0A)) \vee (p \ V1B))) \Rightarrow False) \Leftrightarrow ((p \ V0A) \Rightarrow ((\neg(p \ V1B)) \Rightarrow False)))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p \ V0A)) \Rightarrow False) \Rightarrow (((p \ V0A) \Rightarrow False) \Rightarrow False))) \quad (7)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p \ V0p) \Leftrightarrow (p \ V1q) \Rightarrow (p \ V2r)) \Leftrightarrow (((p \ V0p) \vee (p \ V1q)) \wedge (((p \ V0p) \vee (\neg(p \ V2r))) \wedge (\neg(p \ V1q)) \vee ((p \ V2r) \vee (\neg(p \ V0p)))))))) \quad (8)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((p \ V0p) \Leftrightarrow (\neg(p \ V1q))) \Leftrightarrow (((p \ V0p) \vee (p \ V1q)) \wedge ((\neg(p \ V1q)) \vee (\neg(p \ V0p)))))) \quad (9)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p \ V0p) \Rightarrow (p \ V1q))) \Rightarrow (p \ V0p)))) \quad (10)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p \ V0p) \Rightarrow (p \ V1q))) \Rightarrow (\neg(p \ V1q)))) \quad (11)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p \ V0p) \vee (p \ V1q))) \Rightarrow (\neg(p \ V0p)))) \quad (12)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p \ V0p) \vee (p \ V1q))) \Rightarrow (\neg(p \ V1q)))) \quad (13)$$

Theorem 1

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg(p \ V0p)) \Rightarrow (p \ V1q)) \Rightarrow ((p \ V0p) \vee (p \ V1q))))$$