

thm_2EHolSmt_2EIMP_FALSE
(TMbb6KEVqEoNwsfLzfYMLV1Am3eA6NCNowJ)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 6 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & ((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ & ((\neg False) \Leftrightarrow True))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(\\ & p V0t)))))) \end{aligned} \tag{4}$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t)))\Leftrightarrow(p V0t))) \quad (5)$$

Assume the following.

$$(\forall V0A \in 2.((p V0A)\Rightarrow((\neg(p V0A))\Rightarrow False))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A))\Rightarrow False)\Rightarrow(((p V0A)\Rightarrow False)\Rightarrow False))) \quad (7)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p)\Rightarrow(p V1q)))\Rightarrow(p V0p)))) \quad (8)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p)\Rightarrow(p V1q)))\Rightarrow(\neg(p V1q)))))) \quad (9)$$

Theorem 1 $(\forall V0p \in 2.(((\neg(p V0p))\Rightarrow False)\Rightarrow(p V0p)))$.