

thm_2EHolSmt_2ENOT__FALSE
(TMad6SJPWctofV7DGvT25aiwBSyTx4oWj2x)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 3 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p P \Rightarrow p Q)$ of type ι .

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A_{27a} : \iota. (\lambda V0P \in (2^{A_{27a}}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A_{27a}}))$

Definition 5 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t))$

Definition 6 We define `c_2Ebool_2EF` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$.

Definition 7 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t) \text{ c_2Ebool_2EF } 2))$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((\text{True} \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow \text{True}) \Leftrightarrow \\ & (p V0t)) \wedge (((\text{False} \Leftrightarrow (p V0t)) \Leftrightarrow (\neg (p V0t))) \wedge (((p V0t) \Leftrightarrow \text{False}) \Leftrightarrow (\neg (\\ & p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$(\forall V0t \in 2. ((\neg (\neg (p V0t))) \Leftrightarrow (p V0t))) \tag{3}$$

Assume the following.

$$(\forall V0A \in 2. ((p V0A) \Rightarrow (\neg (p V0A)) \Rightarrow \text{False})) \tag{4}$$

Assume the following.

$$(\forall V0A \in 2. (((\neg (p V0A)) \Rightarrow \text{False}) \Rightarrow (((p V0A) \Rightarrow \text{False}) \Rightarrow \text{False}))) \tag{5}$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p \ V0p) \Rightarrow (p \ V1q))) \Rightarrow (p \ V0p)))) \quad (6)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p \ V0p) \Rightarrow (p \ V1q))) \Rightarrow (\neg(p \ V1q))))) \quad (7)$$

Theorem 1 $(\forall V0p \in 2.((p \ V0p) \Rightarrow ((\neg(p \ V0p)) \Rightarrow False)))$.