

thm_2EHolSmt_2EVALID__IFF__TRUE (TML- wnPxhRiHXnjTnJNcZHwLMXfNah1FsWbh)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_21` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define `c_2Ebool_2E_21` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 7 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 8 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21))$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\ & p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \tag{3}$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \tag{4}$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg(\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))))) \quad (5)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (6)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p))))))) \quad (7)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (8)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))))) \quad (9)$$

Theorem 1 $(\forall V0p \in 2.((p V0p) \Rightarrow ((p V0p) \Leftrightarrow True)))$.