

thm_2EHolSmt_2Ep001 (TMQaNfwH- PzhEvBfZdW4UUduFdHWeJdnusrY)

October 26, 2020

Let $ty_2Ebool_2Eitself : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Ebool_2Eitself\ A0) \quad (1)$$

Let $c_2Ebool_2Ethe_value : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Ebool_2Ethe_value\ A_27a \in (ty_2Ebool_2Eitself\ A_27a) \quad (2)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (3)$$

Let $c_2Ewords_2Edimword : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Ewords_2Edimword\ A_27a \in (ty_2Enum_2Enum^{(ty_2Ebool_2Eitself\ A_27a)}) \quad (4)$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (5)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (6)$$

Definition 1 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$.

Definition 4 We define c_2Ebool_2E21 to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ (ap\ (c_2Emin_2E3D\ (2^{A_27a})))$

Definition 5 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_Emin_E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 7 We define $c_Ebool_E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_Emin_E_3D_3D_3E V0t) c_Ebool_E_7E))$

Definition 8 We define $c_Ebool_E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_Ebool_E_21 2) (\lambda V2t \in 2.$

Let $c_EEnum_E_EREP_num : \iota$ be given. Assume the following.

$$c_EEnum_E_EREP_num \in (\omega^{ty_E_Enum_E_Enum}) \quad (7)$$

Let $c_EEnum_E_ESUC_REP : \iota$ be given. Assume the following.

$$c_EEnum_E_ESUC_REP \in (\omega^{\omega}) \quad (8)$$

Definition 9 We define $c_EEnum_E_ESUC$ to be $\lambda V0m \in ty_E_Enum_E_Enum.(ap c_EEnum_E_EABS_num ($

Definition 10 We define $c_Emin_E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 11 We define $c_Ebool_E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_Emin_E_40$

Definition 12 We define $c_Eprim_rec_E_3C$ to be $\lambda V0m \in ty_E_Enum_E_Enum.\lambda V1n \in ty_E_Enum_E_Enum$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (p (ap (ap c_Eprim_rec_E_3C c_EEnum_E_E0) \\ & (ap (c_Ewords_E_Edimword A_27a) (c_Ebool_E_Ethe_value A_27a)))) \end{aligned} \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (p (ap (ap c_Eprim_rec_E_3C c_EEnum_E_E0) \\ & (ap (c_Ewords_E_Edimword A_27a) (c_Ebool_E_Ethe_value A_27a)))) \end{aligned}$$