

thm_2EHolSmt_2Er006
(TMVcn34i6wDWpJJ7DrYV3RfDBTWYtGYCQe)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A_{27a}}).(ap (ap (c_2Emin_2E_3D (2^{A_{27a}}))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 7 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 8 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\ & p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \tag{3}$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \tag{4}$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p \ V0A) \vee (p \ V1B))) \Rightarrow False) \Leftrightarrow ((p \ V0A) \Rightarrow False) \Rightarrow ((\neg(p \ V1B)) \Rightarrow False)))))) \quad (5)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg(\neg((p \ V0A)) \vee (p \ V1B))) \Rightarrow False) \Leftrightarrow ((p \ V0A) \Rightarrow (\neg(p \ V1B)) \Rightarrow False)))))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p \ V0A)) \Rightarrow False) \Rightarrow (((p \ V0A) \Rightarrow False) \Rightarrow False))) \quad (7)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p \ V0p) \Leftrightarrow (p \ V1q) \Leftrightarrow (p \ V2r)) \Leftrightarrow (((p \ V0p) \vee ((p \ V1q) \vee (p \ V2r))) \wedge (((p \ V0p) \vee (\neg(p \ V2r)) \vee (\neg(p \ V1q)))) \wedge (((p \ V1q) \vee (\neg(p \ V2r)) \vee (\neg(p \ V0p)))) \wedge ((p \ V2r) \vee ((\neg(p \ V1q)) \vee (\neg(p \ V0p)))))))))) \quad (8)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((p \ V0p) \Leftrightarrow (\neg(p \ V1q))) \Leftrightarrow (((p \ V0p) \vee (p \ V1q)) \wedge ((\neg(p \ V1q)) \vee (\neg(p \ V0p)))))) \quad (9)$$

Theorem 1 $(\forall V0p \in 2.((False \Leftrightarrow (p \ V0p)) \Leftrightarrow (\neg(p \ V0p))))$.