

thm_2EHolSmt_2Er012 (TMGdDE- febugAG2hYDwSzky9mEtCzeWsYof7)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2))) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x)$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. \lambda 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 4 We define `c_2Ebool_2EF` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 7 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (the (\lambda x. x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 8 We define `c_2Ebool_2ECOND` to be $\lambda A. \lambda 27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A. 27a. (\lambda V2t2 \in A. 27a. (ap$

Definition 9 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 10 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t)) \text{ c_2Ebool_2E_5C_2F}))$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((\text{True} \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \wedge \text{True}) \Leftrightarrow \\ & (p \ V0t)) \wedge (((\text{False} \wedge (p \ V0t)) \Leftrightarrow \text{False}) \wedge (((p \ V0t) \wedge \text{False}) \Leftrightarrow \text{False}) \wedge \\ & (((p \ V0t) \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((\text{True} \vee (p \ V0t)) \Leftrightarrow \text{True}) \wedge (((p \ V0t) \vee \text{True}) \Leftrightarrow \text{True}) \wedge \\ & (((\text{False} \vee (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \vee \text{False}) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \vee \\ & (p \ V0t)) \Leftrightarrow (p \ V0t)))))) \end{aligned} \tag{3}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (4)$$

Assume the following.

$$(\forall V0b \in 2.(\forall V1t1 \in 2.(\forall V2t2 \in 2.((p (ap (ap (ap (c.2Ebool_2ECOND 2) V0b) V1t1) V2t2)) \Leftrightarrow (((\neg(p V0b)) \vee (p V1t1)) \wedge ((p V0b) \vee (p V2t2)))))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (7)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow (((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (8)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (9)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (10)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ((p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee ((\neg(p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \quad (11)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p)))))) \quad (12)$$

Theorem 1

$$(\forall V0p \in 2.(\forall V1q \in 2.((p (ap (ap (ap (c.2Ebool_2ECOND 2) V0p) V1q) c.2Ebool_2ET)) \Leftrightarrow ((\neg(p V0p)) \vee (p V1q))))$$