

thm\_2EHolSmt\_2Er044  
(TMH3uwRcLKyeH7UHQDXPYZcr6ddWHdFr9g9)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Ebool_2E_21` to be  $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define `c_2Ebool_2E_21` to be  $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A_{27a}}).(ap (ap (c_2Emin_2E_3D (2^{A_{27a}})) (V0P)) (V0P)))$

**Definition 4** We define `c_2Ebool_2E_21` to be  $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow Q)$  of type  $\iota$ .

**Definition 6** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

**Definition 7** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

**Definition 8** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21))$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \tag{3}$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \tag{4}$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (5)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg(\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow (\neg(p V1B) \Rightarrow False)))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (7)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \vee (p V2r)) \Leftrightarrow (((p V0p) \vee (\neg(p V1q))) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((p V1q) \vee ((p V2r) \vee (\neg(p V0p)))))))))) \quad (8)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p)))))) \quad (9)$$

**Theorem 1**  $(\forall V0p \in 2.(((\neg(p V0p)) \vee (p V0p)) \Leftrightarrow True)).$