

thm_2EHolSmt_2Er061
(TMQEcZTWi1cvQXC24wNP61kor7imGEBRjZu)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_7E` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2))) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x)$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A \rightarrow 27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A \rightarrow 27a}))))$

Definition 4 We define `c_2Ebool_2E_7E` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (P \Rightarrow Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t)) (\text{c_2Ebool_2E_7E } V0t)))$

Definition 7 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 8 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. P \text{ (ap } P \text{ x)}) \text{ then } (\lambda x. x \in A \wedge P \text{ x})$ of type $\iota \Rightarrow \iota$.

Definition 9 We define `c_2Ebool_2ECOND` to be $\lambda A. 27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A. 27a. (\lambda V2t2 \in A. 27a. (a$

Definition 10 We define `c_2Ecombin_2EUPDATE` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0a \in A. 27a. \lambda V1b \in A. 27b.$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow (\forall V0x \in A. 27a. ((V0x = V0x) \Leftrightarrow \text{True})) \tag{2}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((\text{True} \Leftrightarrow (p \text{ V0t})) \Leftrightarrow (p \text{ V0t})) \wedge (((p \text{ V0t}) \Leftrightarrow \text{True}) \Leftrightarrow \\ & (p \text{ V0t})) \wedge (((\text{False} \Leftrightarrow (p \text{ V0t})) \Leftrightarrow \neg(p \text{ V0t})) \wedge (((p \text{ V0t}) \Leftrightarrow \text{False}) \Leftrightarrow \neg(\\ & p \text{ V0t)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ & \forall V0f \in (A_27b^{A_27a}). (\forall V1a \in A_27a. ((ap\ (ap\ (ap\ (c_2Ecombin_2EUPDATE \\ & \quad A_27a\ A_27b)\ V1a)\ (ap\ V0f\ V1a))\ V0f) = V0f))) \end{aligned} \tag{4}$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ & \forall V0x \in A_27a. (\forall V1f \in (A_27b^{A_27a}). ((ap\ (ap\ (ap\ (c_2Ecombin_2EUPDATE \\ & \quad A_27a\ A_27b)\ V0x)\ (ap\ V1f\ V0x))\ V1f) = V1f))) \end{aligned}$$