

thm\_2EHolSmt\_2Er254  
(TMGgz9EApqJUBdd7YZYyYPH2MbzT9RoMgby)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Let  $ty\_2Efc\_2Efinite\_image : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Efc\_2Efinite\_image A0) \quad (1)$$

Let  $ty\_2Ebool\_2Eitself : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Ebool\_2Eitself A0) \quad (2)$$

Let  $c\_2Ebool\_2Ethe\_value : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Ebool\_2Ethe\_value A\_27a \in (ty\_2Ebool\_2Eitself A\_27a) \quad (3)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty ty\_2Enum\_2Enum \quad (4)$$

Let  $c\_2Efc\_2Edimindex : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Efc\_2Edimindex A\_27a \in (ty\_2Enum\_2Enum^{(ty\_2Ebool\_2Eitself A\_27a)}) \quad (5)$$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2)) (\lambda V0t \in 2.V0t)$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

**Definition 7** We define  $c\_Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_Ebool\_2E\_21) 2) (\lambda V2t \in 2)))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (6)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (7)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (8)$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num)$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p (ap P x))$  then  $(the (\lambda x.x \in A \wedge P x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40) P)))$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

**Definition 12** We define  $c\_Ebool\_2E\_3F\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap c\_Ebool\_2E\_2F\_5C) P))$

**Definition 13** We define  $c\_2Efcp\_2Efinite\_index$  to be  $\lambda A\_27a : \iota.(ap (c\_2Emin\_2E\_40) (A\_27a^{ty\_2Enum\_2Enum}))$

Let  $ty\_2Efcp\_2Ecart : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Efcp\_2Ecart A0 A1) \quad (9)$$

Let  $c\_2Efcp\_2Edest\_cart : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Efcp\_2Edest\_cart A\_27a A\_27b \in ((A\_27a^{(ty\_2Efcp\_2Efinite\_image A\_27b)})^{(ty\_2Efcp\_2Ecart A\_27a A\_27b)}) \quad (10)$$

**Definition 14** We define  $c\_2Efcp\_2Efcp\_index$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in (ty\_2Efcp\_2Ecart A\_27a A\_27b)$

**Definition 15** We define  $c\_2Efcp\_2EFCP$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.(\lambda V0g \in (A\_27a^{ty\_2Enum\_2Enum}).(ap (c\_2Efcp\_2Efinite\_index) g))$

**Definition 16** We define  $c\_2Ewords\_2Eword\_and$  to be  $\lambda A\_27a : \iota.\lambda V0v \in (ty\_2Efcp\_2Ecart 2 A\_27a).\lambda V0w \in (ty\_2Efcp\_2Ecart 2 A\_27a).$

Assume the following.

$$True \quad (11)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0a \in (ty\_2EfcP\_2Ecart \\ & \quad 2\ A\_27a). (\forall V1b \in (ty\_2EfcP\_2Ecart\ 2\ A\_27a). (\forall V2c \in \\ & \quad (ty\_2EfcP\_2Ecart\ 2\ A\_27a). ((ap\ (ap\ (c\_2Ewords\_2Eword\_and\ A\_27a) \\ & \quad (ap\ (ap\ (c\_2Ewords\_2Eword\_and\ A\_27a)\ V0a)\ V1b))\ V2c) = (ap\ (ap\ ( \\ & \quad c\_2Ewords\_2Eword\_and\ A\_27a)\ V0a)\ (ap\ (ap\ (c\_2Ewords\_2Eword\_and \\ & \quad A\_27a)\ V1b)\ V2c)))))) \end{aligned} \quad (14)$$

**Theorem 1**

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in (ty\_2EfcP\_2Ecart \\ & \quad 2\ A\_27a). (\forall V1y \in (ty\_2EfcP\_2Ecart\ 2\ A\_27a). (\forall V2z \in \\ & \quad (ty\_2EfcP\_2Ecart\ 2\ A\_27a). ((ap\ (ap\ (c\_2Ewords\_2Eword\_and\ A\_27a) \\ & \quad (ap\ (ap\ (c\_2Ewords\_2Eword\_and\ A\_27a)\ V0x)\ V1y))\ V2z) = (ap\ (ap\ ( \\ & \quad c\_2Ewords\_2Eword\_and\ A\_27a)\ V0x)\ (ap\ (ap\ (c\_2Ewords\_2Eword\_and \\ & \quad A\_27a)\ V1y)\ V2z)))))) \end{aligned}$$