

thm\_2ETemporal\_Logic\_2EALWAYS\_IDEM  
 (TMXmvGoiJWPXyCANxZwb-  
 cyRv56SDGwLC1oW)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{2}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2ETemporal\_Logic\_2EALWAYS$  to be  $\lambda V0P \in (2^{ty\_2Enum\_2Enum}).\lambda V1t0 \in ty\_2Enum$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (omega^{ty\_2Enum\_2Enum}) \tag{3}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (omega^{omega}) \tag{4}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{omega}) \tag{5}$$

**Definition 5** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num ($

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in omega \tag{6}$$

**Definition 6** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 7** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o\ (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .

**Definition 8** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in 2.$

**Definition 9** We define  $c\_2Ebool\_2EF$  to be  $(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 10** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_7E$

**Definition 11** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in 2.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & \quad ((ap\ (ap\ c\_2Earithmetic\_2E\_2B\ c\_2Enum\_2E0)\ V0m) = V0m) \wedge (((ap\ ( \\ & \quad ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ c\_2Enum\_2E0) = V0m) \wedge (((ap\ (ap\ c\_2Earithmetic\_2E\_2B \\ & \quad (ap\ c\_2Enum\_2ESUC\ V0m))\ V1n) = (ap\ c\_2Enum\_2ESUC\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B \\ & \quad V0m)\ V1n))) \wedge ((ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ (ap\ c\_2Enum\_2ESUC \\ & \quad V1n)) = (ap\ c\_2Enum\_2ESUC\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n))))))))) \\ & \hspace{15em} (7) \end{aligned}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & \quad \forall V2p \in ty\_2Enum\_2Enum.((ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m) \\ & \quad (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V1n)\ V2p)) = (ap\ (ap\ c\_2Earithmetic\_2E\_2B \\ & \quad (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n))\ V2p)))))) \\ & \hspace{15em} (8) \end{aligned}$$

Assume the following.

$$True \hspace{15em} (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p \\ & \quad V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \\ & \hspace{15em} (10) \end{aligned}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \hspace{15em} (11)$$

Assume the following.

$$(\forall V0t \in 2.((p\ V0t) \vee (\neg(p\ V0t)))) \hspace{15em} (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & \quad True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ & \quad (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t))))))))) \\ & \hspace{15em} (13) \end{aligned}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \hspace{15em} (14)$$

**Theorem 1**

$$(\forall V0a \in (2^{ty\_2Enum\_2Enum}).((ap\ c\_2ETemporal\_Logic\_2EALWAYS \\ V0a) = (ap\ c\_2ETemporal\_Logic\_2EALWAYS (ap\ c\_2ETemporal\_Logic\_2EALWAYS \\ V0a))))$$