

thm_2ETemporal_Logic_2EALWAYS_NEXT (TMH9qLXqcZbEnx2Wwq6o4J4uFEdK2hLnMWe)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 1 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o(x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 3 We define c_2Ebool_2E21 to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E3D\ (2^{A-27a}))\ (\lambda V1P \in 2.V1P))\ (\lambda V1t \in 2.V1t)))$

Definition 4 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (c_2Emin_2E3D\ (2^{ty_2Enum_2Enum}))\ (\lambda V1t \in ty_2Enum_2Enum)))$

Definition 5 We define $c_2ETemporal_Logic_2ENEXT$ to be $\lambda V0P \in (2^{ty_2Enum_2Enum}).(\lambda V1t \in ty_2Enum_2Enum.c_2ESUC\ (ap\ (c_2Emin_2E3D\ (2^{ty_2Enum_2Enum}))\ (\lambda V1t' \in ty_2Enum_2Enum)))$

Let $c_2Earithmetic_2E2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{5}$$

Definition 6 We define $c_2ETemporal_Logic_2EALWAYS$ to be $\lambda V0P \in (2^{ty_2Enum_2Enum}).\lambda V1t0 \in ty_2Enum_2Enum.c_2Ebool_2ET\ (ap\ (c_2Emin_2E3D\ (2^{ty_2Enum_2Enum}))\ (\lambda V1t \in ty_2Enum_2Enum.c_2ESUC\ (ap\ (c_2Emin_2E3D\ (2^{ty_2Enum_2Enum}))\ (\lambda V1t' \in ty_2Enum_2Enum))))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{6}$$

Definition 7 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 8 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow Q)$ of type ι .

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & ((ap\ (ap\ c_2Earithmetic_2E_2B\ c_2Enum_2E0)\ V0m) = V0m) \wedge (((ap\ (\\ & ap\ c_2Earithmetic_2E_2B\ V0m)\ c_2Enum_2E0) = V0m) \wedge (((ap\ (ap\ c_2Earithmetic_2E_2B \\ & (ap\ c_2Enum_2ESUC\ V0m))\ V1n) = (ap\ c_2Enum_2ESUC\ (ap\ (ap\ c_2Earithmetic_2E_2B \\ & V0m)\ V1n))) \wedge ((ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ (ap\ c_2Enum_2ESUC \\ & V1n)) = (ap\ c_2Enum_2ESUC\ (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ V1n))))))))) \\ & \tag{7} \end{aligned}$$

Assume the following.

$$True \tag{8}$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p \\ & V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \\ & \tag{9} \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in \\ & A_27a.(p\ V0t) \Leftrightarrow (p\ V0t))) \\ & \tag{10} \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow \\ & True)) \\ & \tag{11} \end{aligned}$$

Theorem 1

$$\begin{aligned} & (\forall V0a \in (2^{ty_2Enum_2Enum}).((ap\ c_2ETemporal_Logic_2ENEXT \\ & (ap\ c_2ETemporal_Logic_2EALWAYS\ V0a)) = (ap\ c_2ETemporal_Logic_2EALWAYS \\ & (ap\ c_2ETemporal_Logic_2ENEXT\ V0a)))) \end{aligned}$$