

thm_2ETemporal_Logic_2EALWAYS_SIGNAL
(TMH9vMBQB71fXS1obznSaZZwcFYaNh7xMfV)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{2}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2ETemporal_Logic_2EALWAYS$ to be $\lambda V0P \in (2^{ty_2Enum_2Enum}).\lambda V1t0 \in ty_2Enum_2Enum$

Assume the following.

$$True \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{4}$$

Theorem 1

$$(\forall V0a \in (2^{ty_2Enum_2Enum}).(\forall V1t0 \in ty_2Enum_2Enum. ((p (ap (ap c_2ETemporal_Logic_2EALWAYS V0a) V1t0)) \Leftrightarrow (\forall V2t \in ty_2Enum_2Enum.(p (ap V0a (ap (ap c_2Earithmetic_2E_2B V2t) V1t0))))))))$$