

# thm\_2ETemporal\_Logic\_2EBEFORE\_EVENT (TMVomDQJfjgkQjdatXZvpdtHz9pQg6ku8jJ)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num ($

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{5}$$

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

**Definition 7** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 8** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

**Definition 9** We define `c_2ETemporal_Logic_2EWATCH` to be  $\lambda V0q \in (2^{ty\_2Enum\_2Enum}).\lambda V1b \in (2^{ty\_2En$

**Definition 10** We define `c_2Emin_2E_40` to be  $\lambda A.\lambda P \in 2^A$ .**if**  $(\exists x \in A.p (ap P x))$  **then** (the  $(\lambda x.x \in A \wedge$   
of type  $\iota \Rightarrow \iota$ ).

**Definition 11** We define `c_2Ebool_2E_3F` to be  $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A_{27a}}).(ap V0P (ap (c_2Emin_2E_40$

**Definition 12** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E$

**Definition 13** We define `c_2ETemporal_Logic_2EBEFORE` to be  $\lambda V0a \in (2^{ty\_2Enum\_2Enum}).\lambda V1b \in (2^{ty\_2$

Assume the following.

$$True \tag{6}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{7}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{8}$$

Assume the following.

$$\forall A_{27a}.nonempty A_{27a} \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_{27a}.(p V0t)) \Leftrightarrow (p V0t))) \tag{9}$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \tag{10}$$

Assume the following.

$$(\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee (p V0t)) \Leftrightarrow (p V0t)))))) \tag{11}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \tag{12}$$

Assume the following.

$$\forall A_{27a}.nonempty A_{27a} \Rightarrow (\forall V0x \in A_{27a}..((V0x = V0x) \Leftrightarrow True)) \tag{13}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \tag{14}$$

**Theorem 1**

$$\begin{aligned} & (\forall V0a \in (2^{ty\_2Enum\_2Enum}).(\forall V1b \in (2^{ty\_2Enum\_2Enum}). \\ & ((ap (ap c\_2ETemporal\_Logic\_2EBEFORE V0a) V1b) = (ap (ap c\_2ETemporal\_Logic\_2EBEFORE \\ & (\lambda V2t \in ty\_2Enum\_2Enum.(ap (ap c\_2Ebool\_2E\_2F\_5C (ap V0a V2t)) \\ & (ap c\_2Ebool\_2E\_7E (ap V1b V2t)))))) V1b)))) \end{aligned}$$