

thm_2ETemporal_Logic_2EDELTACASES (TMcf4sFKP8KKSEysay7BbKqurLksERuy52A)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum)^{ty_2Enum_2Enum})^{ty_2Enum_2Enum} \tag{2}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o(x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (P)) (V0t) (V1t)))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2)) (\lambda V0t \in 2.V0t)$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o(p \Rightarrow Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.V2t)))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (omega^{ty_2Enum_2Enum}) \tag{3}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (omega^{omega}) \tag{4}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{omega}) \tag{5}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap\ P\ x))$ **then** (the $(\lambda x.x \in A \wedge p$
of type $\iota \Rightarrow \iota$).

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ V0P (ap (c_2Emin_2E_40$

Definition 11 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum$

Definition 12 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21\ 2) (\lambda V2t \in$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty_2Enum_2Enum}).((\exists V1n \in ty_2Enum_2Enum. \\ & (p (ap\ V0P\ V1n))) \Rightarrow (\exists V2n \in ty_2Enum_2Enum.((p (ap\ V0P\ V2n)) \wedge \\ & (\forall V3m \in ty_2Enum_2Enum.((p (ap (ap\ c_2Eprim_rec_2E_3C \\ & V3m)\ V2n)) \Rightarrow (\neg(p (ap\ V0P\ V3m)))))))))) \end{aligned} \quad (6)$$

Assume the following.

$$True \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p \\ & V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \end{aligned} \quad (8)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ & (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((\\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & ((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t)) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ & ((\neg False) \Leftrightarrow True)))) \end{aligned} \quad (13)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\ & p\ V0t)))))) \end{aligned} \quad (15)$$

Theorem 1

$$\begin{aligned} & (\forall V0b \in (2^{ty_2Enum_2Enum}). (\forall V1t0 \in ty_2Enum_2Enum. \\ & ((\exists V2d \in ty_2Enum_2Enum. (\forall V3t \in ty_2Enum_2Enum. \\ & ((p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ V3t)\ V2d)) \Rightarrow (\neg(p\ (ap\ V0b\ (ap\ (ap\ c_2Earithmetic_2E_2B \\ & V3t)\ V1t0)))))) \wedge (p\ (ap\ V0b\ (ap\ (ap\ c_2Earithmetic_2E_2B\ V2d)\ V1t0)))))) \vee \\ & (\forall V4d \in ty_2Enum_2Enum. (\neg(p\ (ap\ V0b\ (ap\ (ap\ c_2Earithmetic_2E_2B \\ & V4d)\ V1t0)))))) \end{aligned}$$