

thm_2ETemporal_Logic_2EEVENTUAL_NEXT (TMVTj9FePD8dgjQTiXH3KqnX4EY4opBhZ1Q)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num ($

Definition 5 We define $c_2ETemporal_Logic_2ENEXT$ to be $\lambda V0P \in (2^{ty_2Enum_2Enum}).(\lambda V1t \in ty_2Enum_2Enum$

Definition 6 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$.

Definition 7 We define $c_2Ebool_2E_3F$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40 A$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{5}$$

Definition 8 We define `c.2ETemporal__Logic_2EEVENTUAL` to be $\lambda V0P \in (2^{ty_2Enum_2Enum}).\lambda V1t0 \in ty.$

Let `c.2Enum_2EZZERO__REP` : ι be given. Assume the following.

$$c.2Enum_2EZZERO__REP \in \omega \quad (6)$$

Definition 9 We define `c.2Enum_2E0` to be $(ap\ c.2Enum_2EABS_num\ c.2Enum_2EZZERO__REP).$

Definition 10 We define `c.2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type $\iota.$

Definition 11 We define `c.2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c.2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & ((ap\ (ap\ c.2Earithmetic_2E_2B\ c.2Enum_2E0)\ V0m) = V0m) \wedge (((ap\ (\\ & ap\ c.2Earithmetic_2E_2B\ V0m)\ c.2Enum_2E0) = V0m) \wedge (((ap\ (ap\ c.2Earithmetic_2E_2B \\ & (ap\ c.2Enum_2ESUC\ V0m))\ V1n) = (ap\ c.2Enum_2ESUC\ (ap\ (ap\ c.2Earithmetic_2E_2B \\ & V0m)\ V1n))) \wedge ((ap\ (ap\ c.2Earithmetic_2E_2B\ V0m)\ (ap\ c.2Enum_2ESUC \\ & V1n)) = (ap\ c.2Enum_2ESUC\ (ap\ (ap\ c.2Earithmetic_2E_2B\ V0m)\ V1n))))))) \end{aligned} \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (9)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A.27a.(p\ V0t) \Leftrightarrow (p\ V0t))) \quad (10)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0x \in A.27a. ((V0x = V0x) \Leftrightarrow True)) \quad (11)$$

Theorem 1

$$\begin{aligned} & (\forall V0a \in (2^{ty_2Enum_2Enum}). ((ap\ c.2ETemporal_Logic_2ENEXT \\ & (ap\ c.2ETemporal_Logic_2EEVENTUAL\ V0a)) = (ap\ c.2ETemporal_Logic_2EEVENTUAL \\ & (ap\ c.2ETemporal_Logic_2ENEXT\ V0a)))) \end{aligned}$$