

thm_2Ealignment_2Ealigned__align (TMHZ5oMuqRFHNYdYfTqggTATzuEqdo9jARc)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define c_2Ebool_2E21 to be $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Let $c_2Earithmetic_2E2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2EBIT1))$

Definition 8 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $ty_2Ebool_2Eitself : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Ebool_2Eitself A0) \quad (7)$$

Let $c_2Ebool_2Ethe_value : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Ebool_2Ethe_value A_27a \in (ty_2Ebool_2Eitself A_27a) \quad (8)$$

Let $c_2Efcp_2Edimindex : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Efcp_2Edimindex A_27a \in (ty_2Enum_2Enum^{(ty_2Ebool_2Eitself A_27a)}) \quad (9)$$

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (10)$$

Let $ty_2Efcp_2Efinite_image : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Efcp_2Efinite_image A0) \quad (11)$$

Definition 9 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2)) (\lambda V0t \in 2.V0t)$.

Definition 10 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 11 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E))$

Definition 12 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.V2t)))$

Definition 13 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then $(the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 14 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a))))$

Definition 15 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum$

Definition 16 We define $c_2Ebool_2E_3F_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap c_2Ebool_2E_2F_5C A_27a P)))$

Definition 17 We define $c_2Efcp_2Efinite_index$ to be $\lambda A_27a : \iota.(ap (c_2Emin_2E_40 (A_27a^{ty_2Enum_2Enum})))$

Let $ty_2Efcp_2Ecart : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Efcp_2Ecart A0 A1) \quad (12)$$

Let $c_2Efcp_2Edest_cart : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Efcp_2Edest_cart A_27a A_27b \in ((A_27a^{(ty_2Efcp_2Efinite_image A_27b)})^{(ty_2Efcp_2Ecart A_27a A_27b)}) \quad (13)$$

Definition 18 We define $c_2Efc_2Efc_index$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in (ty_2Efc_2Ecart\ 2\ A_27a)$

Definition 19 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.$

Definition 20 We define $c_2Earithmetic_2EMIN$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Definition 21 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 22 We define $c_2Earithmetic_2E_3C_2D$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Definition 23 We define c_2Efc_2EFCP to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0g \in (A_27a^{ty_2Enum_2Enum}).(ap$

Definition 24 We define $c_2Ewords_2Eword_slice$ to be $\lambda A_27a : \iota.\lambda V0h \in ty_2Enum_2Enum.\lambda V1l \in ty_2Enum_2Enum.$

Definition 25 We define $c_2Ealignment_2Ealign$ to be $\lambda A_27a : \iota.\lambda V0p \in ty_2Enum_2Enum.\lambda V1w \in (ty_2Enum_2Enum.$

Definition 26 We define $c_2Ealignment_2Ealigned$ to be $\lambda A_27a : \iota.\lambda V0p \in ty_2Enum_2Enum.\lambda V1w \in (ty_2Enum_2Enum.$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0p \in ty_2Enum_2Enum. (\\ & \forall V1w \in (ty_2Efc_2Ecart\ 2\ A_27a). ((ap\ (ap\ (c_2Ealignment_2Ealign \\ & A_27a)\ V0p)\ (ap\ (ap\ (c_2Ealignment_2Ealign\ A_27a)\ V0p)\ V1w))) = (\\ & ap\ (ap\ (c_2Ealignment_2Ealign\ A_27a)\ V0p)\ V1w))) \end{aligned} \quad (14)$$

Assume the following.

$$True \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in \\ & A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow \\ & True)) \end{aligned} \quad (17)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0p \in ty_2Enum_2Enum. (\\ & \forall V1w \in (ty_2Efc_2Ecart\ 2\ A_27a). (p\ (ap\ (ap\ (c_2Ealignment_2Ealigned \\ & A_27a)\ V0p)\ (ap\ (ap\ (c_2Ealignment_2Ealign\ A_27a)\ V0p)\ V1w)))) \end{aligned}$$