

# thm\_2Earithmetic\_2EABS\_\_DIFF\_\_COMM (TM- FuMUAbMd5gByhHBbpjdvXCErPegAZeQC7)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Earithmetic\_2E\_2D : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2D \in ((ty\_2Enum\_2Enum)^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum} \tag{2}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o(x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a})) (P) (V0t) (V1t))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o(p \Rightarrow Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (omega^{ty\_2Enum\_2Enum}) \tag{3}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (omega^{omega}) \tag{4}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{omega}) \tag{5}$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap\ P\ x)) \text{ then } (the\ (\lambda x.x \in A \wedge p$   
of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ V0P (ap (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

**Definition 12** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A\_27a.(\lambda V2t2 \in A\_27a.($

**Definition 13** We define  $c\_2Earithmetic\_2EABS\_DIFF$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.\lambda V1m \in ty\_2Enum$

Assume the following.

$$\begin{aligned} & (\forall V0n \in ty\_2Enum\_2Enum.(\forall V1m \in ty\_2Enum\_2Enum.( \\ & (ap (ap\ c\_2Earithmetic\_2EABS\_DIFF\ V0n)\ V1m) = (ap (ap\ c\_2Earithmetic\_2EABS\_DIFF \\ & \quad V1m)\ V0n)))) \end{aligned} \tag{6}$$

**Theorem 1**

$$\begin{aligned} & (\forall V0n \in ty\_2Enum\_2Enum.(\forall V1m \in ty\_2Enum\_2Enum.( \\ & (ap (ap\ c\_2Earithmetic\_2EABS\_DIFF\ V0n)\ V1m) = (ap (ap\ c\_2Earithmetic\_2EABS\_DIFF \\ & \quad V1m)\ V0n)))) \end{aligned}$$