

thm_2Earithmetic_2EABS__DIFF__EQS (TM- Sqj4eSFgsDE1cHV2D42HUExEVtQhfVDpC)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{4}$$

Definition 3 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ P)\ V0t))$

Definition 5 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow Q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2EF))$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (5)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (6)$$

Definition 9 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap\ c_2Enum_2EABS_num$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A. \lambda P \in 2^A. \mathbf{if} (\exists x \in A. p (ap\ P\ x)) \mathbf{then}$ (the $(\lambda x. x \in A \wedge$
of type $\iota \Rightarrow \iota$).

Definition 11 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P (ap (c_2Emin_2E_40$

Definition 12 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 13 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A_27a. (\lambda V2t2 \in A_27a. ($

Definition 14 We define $c_2Earithmetic_2EABS_DIFF$ to be $\lambda V0n \in ty_2Enum_2Enum. \lambda V1m \in ty_2Enum$

Assume the following.

$$(\forall V0c \in ty_2Enum_2Enum. ((ap (ap\ c_2Earithmetic_2E_2D\ V0c) V0c) = c_2Enum_2E0)) \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (9)$$

Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow (\forall V0b \in 2. (\forall V1t \in A_27a. ((ap (ap (ap (c_2Ebool_2ECOND\ A_27a)\ V0b)\ V1t)\ V1t) = V1t))) \quad (10)$$

Theorem 1

$$(\forall V0n \in ty_2Enum_2Enum. ((ap (ap\ c_2Earithmetic_2EABS_DIFF\ V0n)\ V0n) = c_2Enum_2E0))$$