

thm_2Earithmetic_2EADD1

(TMcaZiYuNw2BriDQdKAowX37TkfKe2iiEfx)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (2)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (3)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (4)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (5)$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x)))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap\ c_2Enum_2EABS_num\ m)$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 7 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earithmetic_2EBIT1\ n)\ V)$

Definition 8 We define $c_2Earthmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum. V0x$.

Definition 9 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 10 We define $c_{\text{CBool_2E_2F_5C}}$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_{\text{CBool_2E_21}} 2) (\lambda V2t \in$

Assume the following.

$$((ap\ c_2Earthmetic_2ENUMERAL\ (ap\ c_2Earthmetic_2EBIT1\ c_2Earthmetic_2EZERO)) = \\ (ap\ c_2Enum_2ESUC\ c_2Enum_2E0)) \quad (7)$$

Assume the following.

Assume the following.

True (9)

Assume the following.

$$\forall A. \exists a. \text{nonempty } A \wedge a \Rightarrow (\forall V0x \in A. \exists a. ((V0x = V0x) \Leftrightarrow \text{True})) \quad (10)$$

Assume the following.

$$(\forall V0P \in (2^{ty_2Enum_2Enum}).(((p (ap V0P c_2Enum_2E0)) \wedge \\ (\forall V1n \in ty_2Enum_2Enum.((p (ap V0P V1n)) \Rightarrow (p (ap V0P (ap c_2Enum_2ESUC \\ V1n))))))) \Rightarrow (\forall V2n \in ty_2Enum_2Enum.(p (ap V0P V2n)))))) \quad (11)$$

Theorem 1

$(\forall V0m \in ty_2Enum_2Enum. ((ap\ c_2Enum_2ESUC\ V0m) = (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT1\ c_2Earithmetic_2EZERO))))))$