

thm\_2Earithmetic\_2EADD\_INV\_0\_EQ  
(TMMafARhNzofob8jr9H9Rmv399Ygj1WBk)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o(x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$ .

**Definition 3** We define  $c\_2Ebool\_2E21$  to be  $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E3D\ (2^{A-27a}))\ (\lambda V1P \in 2.V1P))\ (\lambda V2P \in 2.V2P)))$ .

**Definition 4** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num\ (c\_2Enum\_2EREP\_num\ m))$ .

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{5}$$

**Definition 5** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Earithmetic\_2E2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{6}$$

**Definition 6** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_21\_2$  to be  $(ap (c\_2Ebool\_2E\_21\_2) (\lambda V0t \in 2.V0t))$ .

**Definition 8** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_21\_2))$

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21\_2) (\lambda V2t \in 2.V2t))))$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & ((ap (ap c\_2Earithmetic\_2E\_2B c\_2Enum\_2E0) V0m) = V0m) \wedge (((ap ( \\ & ap c\_2Earithmetic\_2E\_2B V0m) c\_2Enum\_2E0) = V0m) \wedge (((ap (ap c\_2Earithmetic\_2E\_2B \\ & (ap c\_2Enum\_2ESUC V0m)) V1n) = (ap c\_2Enum\_2ESUC (ap (ap c\_2Earithmetic\_2E\_2B \\ & V0m) V1n)))) \wedge ((ap (ap c\_2Earithmetic\_2E\_2B V0m) (ap c\_2Enum\_2ESUC \\ & V1n)) = (ap c\_2Enum\_2ESUC (ap (ap c\_2Earithmetic\_2E\_2B V0m) V1n)))))))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & ((ap (ap c\_2Earithmetic\_2E\_2B V0m) V1n) = V0m) \Rightarrow (V1n = c\_2Enum\_2E0)))) \end{aligned} \quad (8)$$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p \\ & V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (11)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p V0t)))))) \end{aligned} \quad (12)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & ((ap (ap c\_2Earithmetic\_2E\_2B V0m) V1n) = V0m) \Leftrightarrow (V1n = c\_2Enum\_2E0)))) \end{aligned}$$