

thm_2Earithmetic_2EADD__MODULUS__LEFT
(TMbh9dKK8S8B6X2fRnyzJVTkNoXc79xVGJa)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum)^{ty_2Enum_2Enum})^{ty_2Enum_2Enum} \tag{2}$$

Let $c_2Earithmetic_2EMOD : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EMOD \in ((ty_2Enum_2Enum)^{ty_2Enum_2Enum})^{ty_2Enum_2Enum} \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in omega \tag{4}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{omega}) \tag{5}$$

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda \tau a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ P)\ V0t))$

Definition 5 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2EF))$

Definition 8 We define $c_Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_Ebool_2E_21 2) (\lambda V2t \in 2$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (6)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (7)$$

Definition 9 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge$
of type $\iota \Rightarrow \iota$.

Definition 11 We define $c_2Ebool_2E_3F$ to be $\lambda A.\lambda P : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 12 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum$

Assume the following.

$$\begin{aligned} & ((\forall V0n \in ty_2Enum_2Enum.(\forall V1x \in ty_2Enum_2Enum. \\ & ((p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) V0n)) \Rightarrow ((ap (ap c_2Earithmetic_2EMOD \\ & (ap (ap c_2Earithmetic_2E_2B V1x) V0n)) V0n) = (ap (ap c_2Earithmetic_2EMOD \\ & V1x) V0n)))))) \wedge (\forall V2n \in ty_2Enum_2Enum.(\forall V3x \in ty_2Enum_2Enum. \\ & ((p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) V2n)) \Rightarrow ((ap (ap c_2Earithmetic_2EMOD \\ & (ap (ap c_2Earithmetic_2E_2B V2n) V3x)) V2n) = (ap (ap c_2Earithmetic_2EMOD \\ & V3x) V2n)))))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & (\forall V0n \in ty_2Enum_2Enum.(\forall V1x \in ty_2Enum_2Enum.(\\ & (p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) V0n)) \Rightarrow ((ap (ap c_2Earithmetic_2EMOD \\ & (ap (ap c_2Earithmetic_2E_2B V1x) V0n)) V0n) = (ap (ap c_2Earithmetic_2EMOD \\ & V1x) V0n)))) \end{aligned}$$