

thm\_2Earithmetic\_2ECOMPLETE\_INDUCTION  
 (TMbAe-  
 QBCY48dnGnvZ91Sv15mePdFqkZAxEK)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$  **if**  $(\exists x \in A.p\ (ap\ P\ x))$  **then** (the  $(\lambda x.x \in A \wedge p$   
of type  $\iota \Rightarrow \iota$ ).

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty\_2Enum\_2Enum}).((\exists V1n \in ty\_2Enum\_2Enum. \\ & (p\ (ap\ V0P\ V1n))) \Rightarrow (\exists V2n \in ty\_2Enum\_2Enum.((p\ (ap\ V0P\ V2n)) \wedge \\ & (\forall V3m \in ty\_2Enum\_2Enum.((p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C \\ & V3m)\ V2n)) \Rightarrow (\neg(p\ (ap\ V0P\ V3m)))))))))) \end{aligned} \quad (5)$$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p \\ & V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \end{aligned} \quad (7)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (8)$$

Assume the following.

$$\forall A.\lambda 27a.nonempty\ A.\lambda 27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A.\lambda 27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & ((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ & ((\neg False) \Leftrightarrow True))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg \\ & p\ V0t)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0A \in 2.(\forall V1B \in 2.((\neg((p\ V0A) \Rightarrow (p\ V1B))) \Leftrightarrow ((p\ V0A) \wedge \\ & (\neg(p\ V1B)))))) \end{aligned} \quad (13)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0P \in (2^{ty\_2Enum\_2Enum}).((\forall V1n \in ty\_2Enum\_2Enum. \\ & ((\forall V2m \in ty\_2Enum\_2Enum.((p (ap (ap c\_2Eprim\_rec\_2E\_3C \\ V2m) V1n)) \Rightarrow (p (ap V0P V2m)))) \Rightarrow (p (ap V0P V1n)))) \Rightarrow (\forall V3n \in ty\_2Enum\_2Enum. \\ & (p (ap V0P V3n)))))) \end{aligned}$$