

thm_2Earithmetic_2EEQ_ADD_LCANCEL (TMZped374qNUiN8fKzeu8EeCkgWHHjCXBC6)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (1)$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (2)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (3)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (4)$$

Definition 3 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Definition 4 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p \Rightarrow p \ Q)$ of type ι .

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap (ap (c_2Emin_2E_3D (2^{A_27a})) (\lambda V1t \in 2. (ap (c_2Ebool_2E_21 t) (\lambda V2t \in 2. inj_o (V1t = V2t)))))))$

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2. inj_o (V1t2 = V2t))))))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (5)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (6)$$

Definition 7 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ ($

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & ((ap\ (ap\ c_2Earithmetic_2E_2B\ c_2Enum_2E0)\ V0m) = V0m) \wedge (((ap\ (\\ & ap\ c_2Earithmetic_2E_2B\ V0m)\ c_2Enum_2E0) = V0m) \wedge (((ap\ (ap\ c_2Earithmetic_2E_2B\ \\ & (ap\ c_2Enum_2ESUC\ V0m))\ V1n) = (ap\ c_2Enum_2ESUC\ (ap\ (ap\ c_2Earithmetic_2E_2B\ \\ & V0m)\ V1n))) \wedge ((ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ (ap\ c_2Enum_2ESUC\ \\ & V1n)) = (ap\ c_2Enum_2ESUC\ (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ V1n))))))) \\ & \end{aligned} \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in \\ & A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow \quad (10) \\ & True))$$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty_2Enum_2Enum}).(((p\ (ap\ V0P\ c_2Enum_2E0)) \wedge \\ & (\forall V1n \in ty_2Enum_2Enum.((p\ (ap\ V0P\ V1n)) \Rightarrow (p\ (ap\ V0P\ (ap\ c_2Enum_2ESUC\ \\ & V1n))))))) \Rightarrow (\forall V2n \in ty_2Enum_2Enum.(p\ (ap\ V0P\ V2n)))))) \quad (11) \\ & \end{aligned}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & ((ap\ c_2Enum_2ESUC\ V0m) = (ap\ c_2Enum_2ESUC\ V1n)) \Leftrightarrow (V0m = V1n)))) \quad (12) \\ & \end{aligned}$$

Theorem 1

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & \forall V2p \in ty_2Enum_2Enum.(((ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ \\ & V1n) = (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ V2p)) \Leftrightarrow (V1n = V2p)))) \end{aligned}$$