

thm_2Earithmetic_2EEQ_ADD_RCANCEL (TMHW6qrcFWskNJSQUokPJPbebbhHhwB9mFx)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{2}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))))$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2E_2B V0m) V1n) = (ap (ap c_2Earithmetic_2E_2B V1n) V0m)))) \tag{3}$$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\forall V2p \in ty_2Enum_2Enum.(((ap (ap c_2Earithmetic_2E_2B V0m) V1n) = (ap (ap c_2Earithmetic_2E_2B V0m) V2p)) \Leftrightarrow (V1n = V2p)))))) \tag{4}$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\forall V2p \in ty_2Enum_2Enum.(((ap (ap c_2Earithmetic_2E_2B V0m) V1n) = (ap (ap c_2Earithmetic_2E_2B V1n) V2p)) \Leftrightarrow (V0m = V1n))))))$$