

thm_2Earithmetic_2EEQ__LESS__EQ (TMTs- gQjXc5zrYw9z9Rd42JHM4jqWSpJeGWS)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (c_2Enum_2ESUC_REP m))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_2Emin_2E_40$

Definition 11 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 12 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 13 We define $c_2Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum. (p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D\ V0m)\ V0m))) \quad (5)$$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum. (\forall V1m \in ty_2Enum_2Enum. ((p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D\ V0n)\ V1m)) \wedge (p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D\ V1m)\ V0n)))) \Rightarrow (V0n = V1m))) \quad (6)$$

Assume the following.

$$True \quad (7)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (10)$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (V0m = V1n) \Leftrightarrow ((p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D\ V0m)\ V1n)) \wedge (p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D\ V1n)\ V0m))))))$$