

thm_2Earithmetic_2EEQ_MULT_RCANCEL
(TM-
ToCs4gptniSBKNwxhansUEGnwAPvDByeX)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1n \in 2.V1n))\ (\lambda V2t \in 2.V2t))\ (\lambda V3m \in 2.V3m)))$

Definition 6 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))\ (\lambda V3m \in 2.V3m)))$

Let $c_2Earithmetic_2E_2A : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2A \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{4}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & (ap\ (ap\ c_2Earithmetic_2E_2A\ V0m)\ V1n) = (ap\ (ap\ c_2Earithmetic_2E_2A\ V1n)\ V0m)))) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned}
& (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\
& \quad \forall V2p \in ty_2Enum_2Enum. (((ap (ap c_2Earithmetic_2E_2A V0m) \\
V1n) = (ap (ap c_2Earithmetic_2E_2A V0m) V2p)) \Leftrightarrow ((V0m = c_2Enum_2E0) \vee \\
& \quad (V1n = V2p))))))
\end{aligned} \tag{6}$$

Assume the following.

$$True \tag{7}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in \\
& \quad A_27a.(p V0t)) \Leftrightarrow (p V0t)))
\end{aligned} \tag{8}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow \\
& \quad True))
\end{aligned} \tag{9}$$

Theorem 1

$$\begin{aligned}
& (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\
& \quad \forall V2p \in ty_2Enum_2Enum. (((ap (ap c_2Earithmetic_2E_2A V1n) \\
V0m) = (ap (ap c_2Earithmetic_2E_2A V2p) V0m)) \Leftrightarrow ((V0m = c_2Enum_2E0) \vee \\
& \quad (V1n = V2p))))))
\end{aligned}$$