

thm_2Earithmetic_2EEVEN_OR_ODD (TMUq1JNRFyyopPwGgoUDosmJr7fjg43C8vh)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2EODD : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EODD \in (2^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Earithmetic_2EEVEN : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EEVEN \in (2^{ty_2Enum_2Enum}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define $c_2Ebool_2E_F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_F$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum.((p (ap c_2Earithmetic_2EEVEN V0n)) \Leftrightarrow (\neg(p (ap c_2Earithmetic_2EODD V0n)))))) \tag{4}$$

Assume the following.

$$True \quad (5)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (6)$$

Assume the following.

$$(\forall V0t \in 2.(\neg((p\ V0t) \wedge (\neg(p\ V0t)))))) \quad (7)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (9)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p\ V0A) \wedge (p\ V1B))) \Leftrightarrow ((\neg(p\ V0A)) \vee (\neg(p\ V1B)))) \wedge ((\neg((p\ V0A) \vee (p\ V1B))) \Leftrightarrow ((\neg(p\ V0A)) \wedge (\neg(p\ V1B))))))) \quad (10)$$

Theorem 1

$$(\forall V0n \in ty_2Enum_2Enum.((p\ (ap\ c_2Earithmetic_2EVEN\ V0n)) \vee (p\ (ap\ c_2Earithmetic_2ODD\ V0n))))$$