

thm_2Earithmetic_2EEXP__EQ__0 (TMXARHm- pcYRQCe3NW7yUUmXJVvcqG7BhC8Y)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2EBIT1 V0n) c_2Enum_2E0))$

Definition 8 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A.\lambda P : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40 V0P) P)))$

Definition 11 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 12 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 13 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21 2))$

Definition 14 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

Definition 15 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V1n$

Let $c_2Earithmetic_2EEXP : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EEXP \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum})^{ty_2Enum_2Enum} \quad (7)$$

Let $c_2Earithmetic_2E_2A : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2A \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum})^{ty_2Enum_2Enum} \quad (8)$$

Definition 16 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

Assume the following.

$$\begin{aligned} & ((\forall V0m \in ty_2Enum_2Enum.((ap (ap c_2Earithmetic_2EEXP V0m) c_2Enum_2E0) = (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO)))) \wedge (\forall V1m \in ty_2Enum_2Enum.(\forall V2n \in \\ & ty_2Enum_2Enum.((ap (ap c_2Earithmetic_2EEXP V1m) (ap c_2Enum_2ESUC V2n)) = (ap (ap c_2Earithmetic_2E_2A V1m) (ap (ap c_2Earithmetic_2EEXP V1m) V2n)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & ((ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO)) = \\ & (ap c_2Enum_2ESUC c_2Enum_2E0)) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & ((\forall V0m \in ty_2Enum_2Enum.((V0m = c_2Enum_2E0) \vee (\exists V1n \in \\ & ty_2Enum_2Enum.(V0m = (ap c_2Enum_2ESUC V1n)))))) \end{aligned} \quad (11)$$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum. ((\neg(V0n = c_2Enum_2E0)) \Leftrightarrow (p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) V0n)))) \quad (12)$$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\neg((ap (ap c_2Earithmetic_2EEXP (ap c_2Enum_2ESUC V1n)) V0m) = c_2Enum_2E0)))) \quad (13)$$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. ((ap (ap c_2Earithmetic_2E_2A V0m) V1n) = c_2Enum_2E0) \Leftrightarrow ((V0m = c_2Enum_2E0) \vee (V1n = c_2Enum_2E0)))) \quad (14)$$

Assume the following.

$$True \quad (15)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2))))) \quad (16)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \quad (17)$$

Assume the following.

$$(\forall V0t \in 2. (((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))) \quad (18)$$

Assume the following.

$$(\forall V0t \in 2. (((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee (p V0t)) \Leftrightarrow (p V0t)))) \quad (19)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (20)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (21)$$

Assume the following.

$$\forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V0x \in A_{27a}. (\forall V1y \in A_{27a}. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (22)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow \neg(p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p \ V0t)))))) \quad (23)$$

Assume the following.

$$(\forall V0n \in \text{ty_2Enum_2Enum}. (\neg((\text{ap } c_2Enum_2ESUC \ V0n) = c_2Enum_2E0))) \quad (24)$$

Theorem 1

$$(\forall V0n \in \text{ty_2Enum_2Enum}. (\forall V1m \in \text{ty_2Enum_2Enum}. ((\text{ap } (\text{ap } c_2Earithmic_2EEXP \ V0n) \ V1m) = c_2Enum_2E0) \Leftrightarrow ((V0n = c_2Enum_2E0) \wedge (p \ (\text{ap } (\text{ap } c_2Eprim_rec_2E_3C \ c_2Enum_2E0) \ V1m))))))$$