

thm_2Earithmetic_2ELESS__0__CASES (TM- bQBRDoquqiAxjk4pjmgYym7hEybe7hAXZ)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow p \Rightarrow Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 6 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 9 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{5}$$

Definition 10 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 11 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ **if** $(\exists x \in A.p\ (ap\ P\ x))$ **then** *(the $(\lambda x.x \in A \wedge \dots)$ of type $\iota \Rightarrow \iota$).*

Definition 12 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ V0P\ (ap\ (c_2Emin_2E_40$

Definition 13 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum$

Assume the following.

$$True \tag{6}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ & (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \tag{7}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{8}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\ & p\ V0t)))))) \end{aligned} \tag{9}$$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty_2Enum_2Enum}).(((p\ (ap\ V0P\ c_2Enum_2E0)) \wedge \\ & (\forall V1n \in ty_2Enum_2Enum.((p\ (ap\ V0P\ V1n)) \Rightarrow (p\ (ap\ V0P\ (ap\ c_2Enum_2ESUC \\ & V1n)))))) \Rightarrow (\forall V2n \in ty_2Enum_2Enum.(p\ (ap\ V0P\ V2n)))))) \end{aligned} \tag{10}$$

Assume the following.

$$\begin{aligned} & (\forall V0n \in ty_2Enum_2Enum.(p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ c_2Enum_2E0) \\ & (ap\ c_2Enum_2ESUC\ V0n)))) \end{aligned} \tag{11}$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum.((c_2Enum_2E0 = V0m) \vee (p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ c_2Enum_2E0)\ V0m))))$$