

thm\_2Earithmetic\_2ELESS\_\_ADD  
 (TMaUn5FpxRcMd1QG6F4PHdinW7PTDxuEyrf)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (ap P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p \text{ of type } \iota \Rightarrow \iota))$

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y) \text{ of type } \iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A. \lambda a : \iota. (\lambda V0P \in (2^{A-27a}). (ap V0P (ap (c\_2Emin\_2E\_40 A) (c\_2Emin\_2E\_3D (2^A)))$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (2)$$

**Definition 4** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

**Definition 5** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A. \lambda a : \iota. (\lambda V0P \in (2^{A-27a}). (ap (ap (c\_2Emin\_2E\_3D (2^{A-27a})) (c\_2Ebool\_2ET)))$

**Definition 6** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2. V0t))$ .

**Definition 7** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p \Rightarrow p Q) \text{ of type } \iota$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2. (c\_2Ebool\_2ET)))$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in omega \quad (3)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{omega}) \quad (4)$$

**Definition 9** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 10** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E))$

**Definition 11** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (5)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (6)$$

**Definition 12** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2E0\ V0m)$

**Definition 13** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2ESUC\ V0m\ (\lambda V1n \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2E0\ V1n)))$

Assume the following.

$$((\forall V0n \in ty\_2Enum\_2Enum.((ap\ (ap\ c\_2Earithmetic\_2E\_2B\ c\_2Enum\_2E0)\ V0n) = V0n)) \wedge (\forall V1m \in ty\_2Enum\_2Enum.(\forall V2n \in ty\_2Enum\_2Enum.((ap\ (ap\ c\_2Earithmetic\_2E\_2B\ (ap\ c\_2Enum\_2ESUC\ V1m))\ V2n) = (ap\ c\_2Enum\_2ESUC\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V1m)\ V2n))))))) \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (11)$$

Assume the following.

$$\forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

Assume the following.

$$(\forall V0P \in (2^{ty\_2Enum\_2Enum}).(((p (ap V0P c_2Enum\_2E0)) \wedge \\ (\forall V1n \in ty\_2Enum\_2Enum.((p (ap V0P V1n)) \Rightarrow (p (ap V0P (ap c_2Enum\_2ESUC \\ V1n))))))) \Rightarrow (\forall V2n \in ty\_2Enum\_2Enum.(p (ap V0P V2n)))))) \quad (13)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum. (\neg(p (ap (ap c\_2Eprim\_rec\_2E\_3C V0n) c\_2Enum\_2E0)))) \quad (14)$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. (p (ap (ap c_2Eprim_rec_2E_3C V0m) (ap c_2Enum_2ESUC V1n))) \Leftrightarrow ((V0m = V1n) \vee (p (ap (ap c_2Eprim_rec_2E_3C V0m) V1n)))))) \quad (15)$$

### Theorem 1

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. (\\(p \ (ap \ (ap \ c\_2Eprim\_rec\_2E\_3C \ V1n) \ V0m)) \Rightarrow (\exists V2p \in ty\_2Enum\_2Enum. ((ap \ (ap \ c\_2Earithmetic\_2E\_2B \ V2p) \ V1n) = V0m))))))$$