

thm_2Earithmetic_2ELESS__ADD__SUC (TMVydsh12rssF6bbHaoQ4XntzFxRHEQpZtK)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (1)$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (2)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A. \lambda a : \iota. (\lambda V0P \in (2^{A_27a}). (ap (ap (c_2Emin_2E_3D (2^{A_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p \Rightarrow p \ Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2. (ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2. inj_o (V0t1 = V1t2))))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (3)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (4)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (5)$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ m)$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (6)$$

Definition 9 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap\ P\ x)) \text{ then } (\lambda x.x \in A \wedge P(x)) \text{ else } \iota$

Definition 11 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ V0P\ (ap\ (c_2Emin_2E_40\ A_27a)\ V0P)))$

Definition 12 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.(ap\ (c_2Eprim_rec_2E_3C\ V0m)\ V1n)$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.((ap (ap c_2Earithmetic_2E_2B c_2Enum_2E0) V0m) = V0m) \wedge (((ap (ap c_2Earithmetic_2E_2B V0m) c_2Enum_2E0) = V0m) \wedge (((ap (ap c_2Earithmetic_2E_2B (ap c_2Enum_2ESUC V0m)) V1n) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B V0m) V1n))) \wedge ((ap (ap c_2Earithmetic_2E_2B V0m) (ap c_2Enum_2ESUC V1n)) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B V0m) V1n))))))) \\ & \quad (7) \end{aligned}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.((p (ap (ap c_2Eprim_rec_2E_3C (ap c_2Enum_2ESUC V0m)) (ap c_2Enum_2ESUC V1n))) \Leftrightarrow (p (ap (ap c_2Eprim_rec_2E_3C V0m) V1n))))) \\ & \quad (8) \end{aligned}$$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (10)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t))))) \\ & \quad (11) \end{aligned}$$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty_2Enum_2Enum}).(((p\ (ap\ V0P\ c_2Enum_2E0)) \wedge (\forall V1n \in ty_2Enum_2Enum.((p\ (ap\ V0P\ V1n)) \Rightarrow (p\ (ap\ V0P\ (ap\ c_2Enum_2ESUC V1n))))))) \Rightarrow (\forall V2n \in ty_2Enum_2Enum.(p\ (ap\ V0P\ V2n)))) \\ & \quad (12) \end{aligned}$$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum. (p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) (ap c_2Enum_2ESUC V0n)))) \quad (13)$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (p (ap (ap c_2Eprim_rec_2E_3C V0m) (ap (ap c_2Earithmetic_2E_2B V0m) (ap c_2Enum_2ESUC V1n)))))))$$