

thm_2Earithmetic_2ELESS__EQUAL__ANTISYM (TMcz5EBr7yjkSRAEaHyZjCuoE NMicH HuQLN)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 6 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 9 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (ap c_2Enum_2EREP_num (ap c_2Enum_2ESUC_REP m)))$

Definition 10 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.$ **if** $(\exists x \in A.p (ap P x))$ **then** (the $(\lambda x.x \in A \wedge p (ap P x))$ of type $\iota \Rightarrow \iota$).

Definition 11 We define `c_2Ebool_2E_3F` to be $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A_{27a}}). (ap V0P (ap (c_2Emin_2E_40$

Definition 12 We define `c_2Eprim_rec_2E_3C` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Definition 13 We define `c_2Earithmetic_2E_3C_3D` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ \neg & ((p (ap (ap c_2Eprim_rec_2E_3C V0m) V1n)) \wedge (p (ap (ap c_2Eprim_rec_2E_3C \\ & V1n) V0m)))))) \end{aligned} \tag{5}$$

Assume the following.

$$True \tag{6}$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \tag{7}$$

Assume the following.

$$\forall A_{27a}. nonempty A_{27a} \Rightarrow (\forall V0x \in A_{27a}. ((V0x = V0x) \Leftrightarrow True)) \tag{8}$$

Theorem 1

$$\begin{aligned} & (\forall V0n \in ty_2Enum_2Enum. (\forall V1m \in ty_2Enum_2Enum. (\\ ((p (ap (ap c_2Earithmetic_2E_3C_3D V0n) V1m)) \wedge (p (ap (ap c_2Earithmetic_2E_3C_3D \\ & V1m) V0n))) \Rightarrow (V0n = V1m)))) \end{aligned}$$