

thm_2Earithmetic_2ELESS__EQ__LESS__TRANS
 (TMdMnCTKyNzkEiFrpF-
 mUj4uYKAy64ovZv8r)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (ap c_2Enum_2EREP_num (ap c_2Enum_2ESUC_REP V0m)))$

Definition 9 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.$ **if** $(\exists x \in A.p (ap P x))$ **then** (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 10 We define `c_2Ebool_2E_3F` to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 11 We define `c_2Eprim_rec_2E_3C` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum$

Definition 12 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in$

Definition 13 We define `c_2Earithmetic_2E_3C_3D` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & \forall V2p \in ty_2Enum_2Enum.(((p (ap (ap c_2Eprim_rec_2E_3C \\ & V0m) V1n)) \wedge (p (ap (ap c_2Eprim_rec_2E_3C V1n) V2p))) \Rightarrow (p (ap (ap \\ & c_2Eprim_rec_2E_3C V0m) V2p)))))) \end{aligned} \quad (5)$$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p \\ & V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \end{aligned} \quad (7)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2.((p V0t) \vee (\neg (p V0t)))) \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee \\ & (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\ & p\ V0t)))))) \end{aligned} \quad (14)$$

Theorem 1

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & \forall V2p \in ty_2Enum_2Enum. (((p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D \\ & V0m)\ V1n)) \wedge (p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ V1n)\ V2p))) \Rightarrow (p\ (ap\ (ap \\ & c_2Eprim_rec_2E_3C\ V0m)\ V2p)))))) \end{aligned}$$