

thm\_2Earithmetic\_2ELESS\_\_EQ\_\_SUC\_\_REFL  
(TMb6t8rJm4MLtSTKHcZifQeg6cQ7jWcp8Uj)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (2)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (3)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (4)$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num (ap c\_2Enum\_2EREP\_num (ap c\_2Enum\_2ESUC\_REP m)))$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum$

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 13** We define  $c\_2Earithmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum$

Assume the following.

$$True \tag{5}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ & (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p\ V0t)))))) \end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned} & (\forall V0n \in ty\_2Enum\_2Enum. (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ V0n) \\ & (ap\ c\_2Enum\_2ESUC\ V0n)))) \end{aligned} \tag{8}$$

**Theorem 1**

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D \\ & V0m)\ (ap\ c\_2Enum\_2ESUC\ V0m)))) \end{aligned}$$