

# thm\_2Earithmetic\_2ELESS\_LESS\_EQ\_TRANS (TMaPAAYZWhauyR37crHzxjGPrxadrbDaK1)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p \ P \Rightarrow p \ Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2. (ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2. inj\_o (V0t1 = V1t2))))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (2)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (3)$$

Let  $c\_2Enum\_2EAABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EAABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (4)$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. (ap c\_2Enum\_2EAABS\_num m)$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (ap P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 13** We define  $c\_2Earithmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. (\forall V2p \in ty\_2Enum\_2Enum. ((p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ V0m)\ V1n)) \wedge (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ V1n)\ V2p))) \Rightarrow (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ V0m)\ V2p)))))) \quad (5)$$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2))))) \quad (7)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p\ V0t))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2. ((p\ V0t) \vee (\neg(p\ V0t)))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2. (((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \quad (11)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (12)$$

Assume the following.

$$\forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow \text{True})) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2. (((\text{True} \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow \text{True}) \Leftrightarrow (p \ V0t)) \wedge (((\text{False} \Leftrightarrow (p \ V0t)) \Leftrightarrow (\neg(p \ V0t))) \wedge (((p \ V0t) \Leftrightarrow \text{False}) \Leftrightarrow (\neg(p \ V0t))))))) \quad (14)$$

### Theorem 1

$$(\forall V0m \in \text{ty\_2Enum\_2Enum}. (\forall V1n \in \text{ty\_2Enum\_2Enum}. (\forall V2p \in \text{ty\_2Enum\_2Enum}. (((p \ (\text{ap} \ (ap \ c\_2Eprim\_rec\_2E\_3C \ V0m) \ V1n)) \wedge (p \ (\text{ap} \ (ap \ c\_2Earithmetic\_2E\_3C\_3D \ V1n) \ V2p))) \Rightarrow (p \ (\text{ap} \ (ap \ c\_2Eprim\_rec\_2E\_3C \ V0m) \ V2p)))))))$$