

thm\_2Earithmetic\_2ELESS\_\_MONO\_\_ADD\_\_INV  
(TMFnyMg-  
PQXwVy5jCA1j3Rv2iW44S3g5pg3N)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum)^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum} \tag{2}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_7E$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{3}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{4}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{5}$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap\ P\ x))$  **then** (the  $(\lambda x.x \in A \wedge p$   
of type  $\iota \Rightarrow \iota$ ).

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ V0P (ap (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{6}$$

**Definition 12** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & ((ap (ap\ c\_2Earithmetic\_2E\_2B\ c\_2Enum\_2E0)\ V0m) = V0m) \wedge (((ap ( \\ & ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ c\_2Enum\_2E0) = V0m) \wedge (((ap (ap\ c\_2Earithmetic\_2E\_2B \\ & (ap\ c\_2Enum\_2ESUC\ V0m))\ V1n) = (ap\ c\_2Enum\_2ESUC (ap (ap\ c\_2Earithmetic\_2E\_2B \\ & V0m)\ V1n))) \wedge ((ap (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ (ap\ c\_2Enum\_2ESUC \\ & V1n)) = (ap\ c\_2Enum\_2ESUC (ap (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n))))))))) \end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & (p (ap (ap\ c\_2Eprim\_rec\_2E\_3C (ap\ c\_2Enum\_2ESUC\ V0m)) (ap\ c\_2Enum\_2ESUC \\ & V1n))) \Leftrightarrow (p (ap (ap\ c\_2Eprim\_rec\_2E\_3C\ V0m)\ V1n)))))) \end{aligned} \tag{8}$$

Assume the following.

$$True \tag{9}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ & (p\ V0t) \Rightarrow False) \Leftrightarrow \neg (p\ V0t)))))) \end{aligned} \tag{10}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg ( \\ & p\ V0t)))))) \end{aligned} \tag{11}$$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty\_2Enum\_2Enum}).(((p (ap\ V0P\ c\_2Enum\_2E0)) \wedge \\ & (\forall V1n \in ty\_2Enum\_2Enum.((p (ap\ V0P\ V1n)) \Rightarrow (p (ap\ V0P (ap\ c\_2Enum\_2ESUC \\ & V1n)))))) \Rightarrow (\forall V2n \in ty\_2Enum\_2Enum.(p (ap\ V0P\ V2n)))))) \end{aligned} \tag{12}$$

**Theorem 1**

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. (\forall V2p \in ty\_2Enum\_2Enum. ((p (ap (ap c\_2Eprim\_rec\_2E\_3C (ap (ap c\_2Earithmetic\_2E\_2B V0m) V2p)) (ap (ap c\_2Earithmetic\_2E\_2B V1n) V2p)))) \Rightarrow (p (ap (ap c\_2Eprim\_rec\_2E\_3C V0m) V1n))))))$$