

thm\_2Arithmetic\_2ELESS\_\_MONO\_\_EQ  
 (TMQocZ-  
 cAVmN7FfQQdzyxzY8Q1dZ13YgzDjq)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num ($

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2$

**Definition 9** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p \text{ (ap } P \ x)) \text{ then (the } (\lambda x. x \in A \wedge p \text{ of type } \iota \Rightarrow \iota).$

**Definition 10** We define `c_2Ebool_2E_3F` to be  $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P \text{ (ap (c_2Emin_2E_40$

**Definition 11** We define `c_2Eprim_rec_2E_3C` to be  $\lambda V0m \in \text{ty\_2Enum\_2Enum}. \lambda V1n \in \text{ty\_2Enum\_2Enum}.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in \text{ty\_2Enum\_2Enum}. (\forall V1n \in \text{ty\_2Enum\_2Enum}. \\ & (p \text{ (ap (ap c_2Eprim\_rec\_2E\_3C (ap c_2Enum\_2ESUC } V0m)) \text{ (ap c_2Enum\_2ESUC } \\ & \quad V1n)))) \Leftrightarrow (p \text{ (ap (ap c_2Eprim\_rec\_2E\_3C } V0m) \text{ } V1n)))) \end{aligned} \quad (5)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0m \in \text{ty\_2Enum\_2Enum}. (\forall V1n \in \text{ty\_2Enum\_2Enum}. \\ & (p \text{ (ap (ap c_2Eprim\_rec\_2E\_3C (ap c_2Enum\_2ESUC } V0m)) \text{ (ap c_2Enum\_2ESUC } \\ & \quad V1n)))) \Leftrightarrow (p \text{ (ap (ap c_2Eprim\_rec\_2E\_3C } V0m) \text{ } V1n)))) \end{aligned}$$