

thm\_2Earithmetic\_2ELESS\_OR  
(TMZKr2ZuuUdYr9JGwA9eoQeDCRqoPoQFxrX)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num ($

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  then (the  $(\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ ).

**Definition 10** We define  $c\_Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap\ V0P\ (ap\ (c\_Emin\_2E\_40$

**Definition 11** We define  $c\_Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 12** We define  $c\_Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c\_Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 13** We define  $c\_Earithmic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\ & (p\ (ap\ (ap\ c\_Eprim\_rec\_2E\_3C\ V0m)\ V1n)) \Leftrightarrow (p\ (ap\ (ap\ c\_Earithmic\_2E\_3C\_3D \\ & (ap\ c\_Eenum\_2ESUC\ V0m))\ V1n)))))) \end{aligned} \quad (5)$$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t))))) \end{aligned} \quad (8)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\ & (p\ (ap\ (ap\ c\_Eprim\_rec\_2E\_3C\ V0m)\ V1n)) \Rightarrow (p\ (ap\ (ap\ c\_Earithmic\_2E\_3C\_3D \\ & (ap\ c\_Eenum\_2ESUC\ V0m))\ V1n)))))) \end{aligned}$$