

thm_2Earithmetic_2ELESS_OR_EQ_ALT
(TMF717qqrGACbYk4LAcBLokSwo5RB6cinom)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (c_2Enum_2ESUC_REP m))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define `c_2Ebool_2E_3F` to be $\lambda A_{.27a} : \iota. (\lambda V0P \in (2^{A_{.27a}}). (ap\ V0P\ (ap\ (c_2Emin_2E_40$

Definition 11 We define `c_2Eprim_rec_2E_3C` to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 12 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 13 We define `c_2Earithmetic_2E_3C_3D` to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 14 We define `c_2ERelation_2ETC` to be $\lambda A_{.27a} : \iota. \lambda V0R \in ((2^{A_{.27a}})^{A_{.27a}}). \lambda V1a \in A_{.27a}. \lambda V2b \in$

Definition 15 We define `c_2ERelation_2ERTC` to be $\lambda A_{.27a} : \iota. \lambda V0R \in ((2^{A_{.27a}})^{A_{.27a}}). \lambda V1a \in A_{.27a}. \lambda V2b \in$

Assume the following.

$$True \tag{5}$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \tag{6}$$

Assume the following.

$$(\forall V0t \in 2. (((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \tag{7}$$

Assume the following.

$$\forall A_{.27a}. nonempty\ A_{.27a} \Rightarrow (\forall V0x \in A_{.27a}. ((V0x = V0x) \Leftrightarrow True)) \tag{8}$$

Assume the following.

$$\forall A_{.27a}. nonempty\ A_{.27a} \Rightarrow \forall A_{.27b}. nonempty\ A_{.27b} \Rightarrow (\forall V0f \in (A_{.27b}^{A_{.27a}}). (\forall V1g \in (A_{.27b}^{A_{.27a}}). ((V0f = V1g) \Leftrightarrow (\forall V2x \in A_{.27a}. ((ap\ V0f\ V2x) = (ap\ V1g\ V2x)))))) \tag{9}$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \tag{10}$$

Assume the following.

$$(c_2Eprim_rec_2E_3C = (ap\ (c_2ERelation_2ETC\ ty_2Enum_2Enum)\ (\lambda V0x \in ty_2Enum_2Enum. (\lambda V1y \in ty_2Enum_2Enum. (ap\ (ap\ (c_2Emin_2E_3D\ ty_2Enum_2Enum)\ V1y)\ (ap\ c_2Enum_2ESUC\ V0x)))))) \tag{11}$$

Assume the following.

$$\begin{aligned}
& \forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0R \in ((2^{A.27a})^{A.27a}). \\
& (\forall V1x \in A.27a. (\forall V2y \in A.27a. ((p\ (ap\ (ap\ (ap\ (c.2Erelation.2ERTC \\
& A.27a)\ V0R)\ V1x)\ V2y))) \Leftrightarrow ((V1x = V2y) \vee (p\ (ap\ (ap\ (ap\ (c.2Erelation.2ETC \\
& A.27a)\ V0R)\ V1x)\ V2y))))))
\end{aligned} \tag{12}$$

Theorem 1

$$\begin{aligned}
(c.2Earithmic.2E.3C.3D = (ap\ (c.2Erelation.2ERTC\ ty.2Enum.2Enum) \\
(\lambda V0x \in ty.2Enum.2Enum. (\lambda V1y \in ty.2Enum.2Enum. (ap\ (ap\ (\\
c.2Emin.2E.3D\ ty.2Enum.2Enum)\ V1y)\ (ap\ c.2Enum.2ESUC\ V0x))))))
\end{aligned}$$