

thm_2Earithmetic_2ELESS__STRONG__ADD
 (TMHTupCehZoKGThmbhks5q6TjkJy7goDqd9S)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (2)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (3)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p \Rightarrow p\ Q)$ of type ι .

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2. V0x))\ (\lambda V1x \in 2. V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ (ap\ (c_2Emin_2E_3D\ (2^{A_27a}))\ (\lambda V1t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2. inj_o (V2t = t1)))))))$

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2. inj_o (V2t = t1))))))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (4)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (5)$$

Definition 7 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap\ c_2Enum_2EABS_num\ m)$

Definition 8 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 9 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF))$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (\text{the } (\lambda x.x \in A \wedge p \text{ of type } \iota \Rightarrow \iota))$

Definition 11 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 12 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 13 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in$

Definition 14 We define $c_2Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & ((ap (ap c_2Earithmetic_2E_2B c_2Enum_2E0) V0m) = V0m) \wedge (((ap (\\ & ap (ap c_2Earithmetic_2E_2B V0m) c_2Enum_2E0) = V0m) \wedge (((ap (ap c_2Earithmetic_2E_2B \\ & (ap c_2Enum_2ESUC V0m)) V1n) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B \\ & V0m) V1n))) \wedge ((ap (ap c_2Earithmetic_2E_2B V0m) (ap c_2Enum_2ESUC \\ & V1n)) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B V0m) V1n))))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & (p (ap (ap c_2Eprim_rec_2E_3C V0m) V1n)) \Rightarrow (p (ap (ap c_2Earithmetic_2E_3C_3D \\ & (ap c_2Enum_2ESUC V0m)) V1n)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & (p (ap (ap c_2Earithmetic_2E_3C_3D V1n) V0m)) \Rightarrow (\exists V2p \in ty_2Enum_2Enum. \\ & ((ap (ap c_2Earithmetic_2E_2B V2p) V1n) = V0m)))) \end{aligned} \quad (9)$$

Assume the following.

$$True \quad (10)$$

Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (11)$$

Theorem 1

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & (p (ap (ap c_2Eprim_rec_2E_3C V1n) V0m)) \Rightarrow (\exists V2p \in ty_2Enum_2Enum. \\ & ((ap (ap c_2Earithmetic_2E_2B (ap c_2Enum_2ESUC V2p)) V1n) = V0m)))) \end{aligned}$$