

thm_2Earithmetic_2EMODEQ__0__CONG
(TMPybtnhEfTrGrstt1JqKAx2LU8T7LhBaYL)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o(x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(\lambda p.\lambda q.(c_2Emin_2E_3D(2^2))(\lambda V0x \in 2.V0x))(\lambda V1x \in 2.V1x)$.

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(\lambda p.\lambda q.(c_2Emin_2E_3D(2^{A-27a}))(\lambda V0x \in 2.V0x))(\lambda V1x \in 2.V1x)))$.

Definition 4 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(\lambda p.\lambda q.(c_2Enum_2EABS_num(p)(q)))$.

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{5}$$

Definition 5 We define c_2Enum_2E0 to be $(\lambda p.\lambda q.(c_2Enum_2EABS_num(p)(q)(c_2Enum_2EZERO_REP)))$.

Definition 6 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2EBIT1 V0n) V0n)$

Definition 8 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 9 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 10 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

Let $c_2Earithmetic_2E_2A : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2A \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (7)$$

Definition 11 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge P x)$) of type $\iota \Rightarrow \iota$.

Definition 12 We define $c_2Ebool_2E_3F$ to be $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^A)^{27a}).(ap V0P (ap (c_2Emin_2E_40 V0P)))$

Definition 13 We define $c_2Earithmetic_2EMODEQ$ to be $\lambda V0n \in ty_2Enum_2Enum.\lambda V1m1 \in ty_2Enum_2Enum.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & ((ap (ap c_2Earithmetic_2E_2B c_2Enum_2E0) V0m) = V0m) \wedge (((ap (\\ & ap c_2Earithmetic_2E_2B V0m) c_2Enum_2E0) = V0m) \wedge (((ap (ap c_2Earithmetic_2E_2B \\ & (ap c_2Enum_2ESUC V0m)) V1n) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B \\ & V0m) V1n))) \wedge ((ap (ap c_2Earithmetic_2E_2B V0m) (ap c_2Enum_2ESUC \\ & V1n)) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B V0m) V1n)))))))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & ((ap (ap c_2Earithmetic_2E_2A c_2Enum_2E0) V0m) = c_2Enum_2E0) \wedge \\ & (((ap (ap c_2Earithmetic_2E_2A V0m) c_2Enum_2E0) = c_2Enum_2E0) \wedge \\ & (((ap (ap c_2Earithmetic_2E_2A (ap c_2Earithmetic_2ENUMERAL \\ & (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO))) V0m) = V0m) \wedge \\ & (((ap (ap c_2Earithmetic_2E_2A V0m) (ap c_2Earithmetic_2ENUMERAL \\ & (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO))) = V0m) \wedge (\\ & ((ap (ap c_2Earithmetic_2E_2A (ap c_2Enum_2ESUC V0m)) V1n) = (ap \\ & (ap c_2Earithmetic_2E_2B (ap (ap c_2Earithmetic_2E_2A V0m) V1n)) \\ & V1n)) \wedge ((ap (ap c_2Earithmetic_2E_2A V0m) (ap c_2Enum_2ESUC V1n)) = \\ & (ap (ap c_2Earithmetic_2E_2B V0m) (ap (ap c_2Earithmetic_2E_2A \\ & V0m) V1n)))))))))) \end{aligned} \quad (9)$$

Assume the following.

$$True \quad (10)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\exists V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (11)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

Theorem 1

$$(\forall V0m1 \in ty_2Enum_2Enum. (\forall V1m2 \in ty_2Enum_2Enum. ((p\ (ap\ (ap\ (ap\ c_2Earithmetic_2EMODEQ\ c_2Enum_2E0)\ V0m1)\ V1m2)) \Leftrightarrow (V0m1 = V1m2))))$$